



KEMENTERIAN KESIHATAN MALAYSIA



KEMENTERIAN PENDIDIKAN MALAYSIA

## NATIONAL HEALTH & MORBIDITY SURVEY 2022

# ADOLESCENT HEALTH SURVEY 2022



**WP LABUAN**



**Contributors**

The following persons contributed to the interpretation of findings, discussions on implications, conclusions and/or drawing recommendations for this report.

**(In alphabetical order)**

Ahmad Ali Zainuddin, Ainan Nasrina Ismail, Anita Suleiman, Annapurny Venkiteswaran, Azli Baharudin@Shaharudin, Chan Yee Mang, Chong Chean Tat, Chong Zhuo Lin, Eida Nurhadzira Muhammad, Faizul Akmal Abdul Rahim, Fatin Athira Tahir, Fazila Haryati Ahmad, Filza Noor Asari, Halizah Mat Rifin, Hamizatul Akmal Abd Hamid, Hasimah Ismail, Hazizi Abu Saad, Khairul Hasnan Amali, Khaw Wan-Fei, Kishwen Kanna Yoga Ratnam, Lai Wai Kent, Lalitha Palaniveloo, Liew Siaw Hun, Lim Kuang Kuay, Maznieda Mahjom, Mohamad Aznuddin Abd Razak, Mohamad Salleh Abdul Ghani, Mohd Amierul Fikri Mahmud, Mohd Azahadi Omar, Mohd Farihan Md Yatim, Mohd Haniff Bistari, Mohd Hatta Abdul Mutalip, Mohd Hairmanshah Mohd Shah, Mohd Hazrin Hasim@Hashim, Mohd Ruhaizie Riyadzi, Mohd Shaiful Azlan Kassim, Muhamad Khairul Nazrin Khalil, Muhammad Azri Adam Adnan, Muhammad Fadhli Mohd Yusoff, Muhammad Faiz Mohd Hashim, Muhammad Hanafi Bakri, Muhammad Solihin Rezali, Munawara Pardi, Murnizar Mokhtar, Musalnizan Mustalkah, Nazirah Alias, Nik Adilah Shahein, Nik Daliana Nik Farid, Nik Rubiah Nik Abdul Rashid, Nik Ruzyanei Nik Jaafar, Nizam Baharom, Noor Aliza Lodz, Noor Raihan Khamal, Noor Suraya Muhamad, Noor Syaqlah Shawaluddin, Nor Rahidah Abd Rahim, Nor'Ain Ab Wahab, Noraryana Hassan, Norhafizah Sahril, Norhayati Nordin, Norlaila Hamid, Norli Abd Jabbar, Norliana Ismail, Norliza Shamsuddin, Norsyamalina Che Abdul Rahim, Norzawati Yoep, Nur Faraeein Zainal Abidin, Nur Hamizah Nasaruddin, Nur Hidayatun Fadhilah Mohd Nor, Nurashikin Ibrahim, Nurul Haniyah Rosslan, Nurul Huda Ibrahim, Nurulasmak Mohamed, Nurzaime Zulaily, Rafidah Ali, Rusdi Abd Rashid, S Maria Awaluddin, Saidatul Norbaya Buang, Sheikh Shafizal Sheikh Ilman, Sherina Mohd Sidek, Shubash Shander Ganapathy, Siti Balkhis Shafie, Siti Adibah Ab Halim, Suhaila Abd Ghaffar, Sulhariza Husni Zain, Syafinaz Mohd Sallehuddin, Tan Lee Ann, Tania Gayle Robert Lourdes, Teh Wai Siew, Thamil Arasu Saminathan, Tuan Amin Tuan Lah, Ummi Nadiyah Yusoff, Wan Kim Sui, Wan Sarifah Aini Wan Jusoh, Zamzaireen Zainal Abidin, Zulkarnain Ramli.

**Editorial Reviewers**

Khaw Wan-Fei, Mohd Hatta Abdul Mutalip, Muhammad Solihin Rezali, Tania Gayle Robert Lourdes, Thamil Arasu Saminathan

**Editors**

Lim Kuang Kuay, Noor Ani Ahmad

**External Reviewer**

Saidatul Norbaya Buang  
Family Health Development Division, Ministry of Health Malaysia

© 2022, Institute for Public Health, National Institutes of Health, Ministry of Health Malaysia, Kuala Lumpur.

Perpustakaan Negara Malaysia

Cataloguing-in-Publication Data

National Health and Morbidity Survey 2022 (NMRR-21-157-58261)  
Malaysia Adolescent Health Survey 2022  
ISBN 978-967-5340-75-8

**MOH/S/IKU/219.23(BK)**

**Suggested citation:**

Institute for Public Health (IPH) 2022. Technical Report National Health and Morbidity Survey (NHMS) 2022: Adolescent Health Survey, WP Labuan.

**Produced and Distributed by:**

National Health and Morbidity Survey 2022: Adolescent Health Survey  
Institute for Public Health  
National Institutes of Health  
Ministry of Health Malaysia  
Blok B5 & B6, Kompleks NIH,  
No1, Jalan Setia Murni U13/52,  
Seksyen U13 Bandar Setia Alam,  
40170 Shah Alam, Selangor.

Tel: +603-3362-7800 Fax: +603-3362-7801

Any inquiries or comments on this report should be directed to the following:

Principal Investigator  
National Health and Morbidity Survey 2022: Adolescent Health Survey  
Institute for Public Health  
National Institutes of Health  
Ministry of Health Malaysia  
Blok B5 & B6, Kompleks NIH,  
No1, Jalan Setia Murni U13/52,  
Seksyen U13 Bandar Setia Alam,  
40170 Shah Alam, Selangor.

Tel: +603-3362-7800 Fax: +603-3362-7801

Published by Institute for Public Health, National Institutes of Health, Ministry of Health Malaysia.



## ACKNOWLEDGEMENT

---

The authors would like to thank the Director General of Health Malaysia and the National Health and Morbidity Survey (NHMS) Steering Committee Team for supporting this research project via financial grant and technical support. Special thanks to the Deputy Director General of Health (Research and Technical Support) and the Director of Institute for Public Health for their continuous advice, guidance and support throughout the study. Gratitude to the Director General of Education, Ministry of Education Malaysia, various State Education Departments, District Education Offices, School Principals and the selected school officers who had been very supportive during the preparation and implementation of data collection. Thanks to all field supervisors, data collectors and individuals who had assisted in the conduct of the study. And last but not least, our sincere appreciation to all students from 239 schools who participated in this study.



## LIST OF ABBREVIATIONS

---

AHS	Adolescent Health Survey
BOD	Burden of Disease
CDC	Centers for Disease Control and Prevention
GSHS	Global School-based Student Health survey
IPH	Institute for Public Health
NMRR	National Medical Research Register
UNICEF	United Nations Children’s Fund
UNESCO	United Nations Educational, Scientific and Cultural Organization
UNODC	United Nation Office on Drug and Crime
WHO	World Health Organization

## TABLE OF CONTENTS

<b>1.0</b>	<b>INTRODUCTION</b>	<b>1</b>
1.1	Objectives	1
1.1.1	General Objectives	1
1.1.2	Specific Objectives	1
<b>2.0</b>	<b>METHODOLOGY</b>	<b>2</b>
2.1	Study Design	2
2.2	Sampling Frame and Target Population	2
2.3	Sample Size Calculation	2
2.4	Sampling Design	2
2.5	Ethical Approval and Consent Forms	2
2.6	Study Instrument	3
2.7	Data Collection	3
2.8	Quality Control	3
2.9	Data Processing and Quality Centre	3
2.10	Data Analysis	3
2.11	References	3
<b>3.0</b>	<b>FINDINGS</b>	<b>6</b>
3.1	General Findings	6
3.2	Alcohol Consumption	8
3.2.1	Introduction	8
3.2.2	Objectives	8
3.2.3	Variable definitions	8
3.2.4	Findings	8
3.2.5	Discussion / Conclusion	8
3.2.6	Recommendations	8
3.2.7	References	8
3.3	Dietary Behaviours	10
3.3.1	Introduction	10
3.3.2	Objectives	10
3.3.3	Variable definitions	10
3.3.4	Findings	10
3.3.5	Discussions / Conclusion	11
3.3.6	Recommendations	11
3.3.7	References	11
3.4	Nutritional Status	17
3.4.1	Introduction	17
3.4.2	Objectives	17
3.4.3	Variable definitions	17
3.4.4	Findings	17
3.4.5	Discussion / Conclusion	17
3.4.6	Recommendations	17
3.4.7	References	17
3.5	Drug Use	20
3.5.1	Introduction	20
3.5.2	Objectives	20
3.5.3	Variable Definitions	20
3.5.4	Findings	20
3.5.5	Discussion / Conclusion	20
3.5.6	Recommendations	21
3.5.7	References	21

## TABLE OF CONTENTS

3.6	Oral and Hand Hygiene	25
3.6.1	Introduction	25
3.6.2	Objectives	25
3.6.3	Variable definitions	25
3.6.4	Findings	25
3.6.5	Discussion / Conclusion	25
3.6.6	Recommendations	26
3.6.7	References	26
3.7	Mental Health Problems	33
3.7.1	Mental Health Problems	33
3.7.1.1	Introduction	33
3.7.1.2	Objectives	33
3.7.1.3	Variable definitions	33
3.7.1.4	Findings	33
3.7.1.5	Discussion / Conclusion	33
3.7.1.6	Recommendations	33
3.7.1.7	References	34
3.7.2	Depression	35
3.7.2.1	Introduction	35
3.7.2.2	Objectives	35
3.7.2.3	Variable definitions	35
3.7.2.4	Findings	35
3.7.2.5	Discussion / Conclusion	35
3.7.2.6	Recommendations	35
3.7.2.7	References	35
3.8	Physical Activity	40
3.8.1	Introduction	40
3.8.2	Objectives	40
3.8.3	Variable Definitions	40
3.8.4	Findings	40
3.8.5	Discussion / Conclusion	40
3.8.6	Recommendations	40
3.8.7	References	41
3.9	Protective Factors	44
3.9.1	Introduction	44
3.9.2	Objectives	44
3.9.3	Variable definitions	44
3.9.4	Findings	44
3.9.5	Discussion / Conclusion	45
3.9.6	Recommendations	45
3.9.7	References	45
3.10	Sexual Behaviours	49
3.10.1	Introduction	49
3.10.2	Objectives	49
3.10.3	Variable Definitions	49
3.10.4	Findings	49
3.10.5	Discussion / Conclusion	50
3.10.6	Recommendations	50
3.10.7	References	50



TABLE OF CONTENTS

3.11	Tobacco Use	53
3.11.1	Introduction	53
3.11.2	Objectives	53
3.11.3	Variable Definitions	80
3.11.4	Findings	53
3.11.5	Discussion / Conclusion	53
3.11.6	Recommendation	53
3.11.7	References	54
3.12	Violence and Unintentional Injury	57
3.12.1	Introduction	57
3.12.2	Objectives	57
3.12.3	Variable Definitions	57
3.12.4	Findings	57
3.12.5	Discussion / Conclusion	58
3.12.6	Recommendations	58
3.12.7	References	58
3.13	Adolescents' Perspectives on the Impact of COVID-19 on their families	64
3.13.1	Introduction	64
3.13.2	Objectives	64
3.13.3	Variable definitions	64
3.13.4	Findings	64
3.13.5	Discussion / Conclusion	64
3.13.6	Recommendations	64
3.13.7	References	64
	<b>APPENDICES</b>	<b>66</b>

## LIST OF TABLES

### METHODOLOGY

**Table 2.1** Distribution of secondary schools sampled, by State

### GENERAL FINDINGS

**Table 3.1.1** Response rate at student level, by state, 2022

### ALCOHOL CONSUMPTION

**Table 3.2.1** Prevalence of ever alcohol drinkers among adolescents in WP Labuan, 2022

**Table 3.2.2** Proportion of ever alcohol drinkers according to alcohol initiation age among adolescents in WP Labuan, 2022

**Table 3.2.3** Usual sources of obtaining alcohol in the past 30 days among current drinkers among adolescents in WP Labuan, 2022

**Table 3.2.4** Number of times (got into trouble with family or friends, missed school or got into a fight as a result of drinking alcohol) among ever alcohol drinkers (proportion) among adolescents in WP Labuan, 2022

### DIETARY BEHAVIOURS

**Table 3.3.1** Prevalence of adolescents in WP Labuan who most of the time or always went hungry in the past 30 days because there was not enough food in his/her home, 2022

**Table 3.3.2** Prevalence of fruit intake of at least twice daily in the past 30 days among adolescents in WP Labuan, 2022

**Table 3.3.3** Prevalence of vegetables intake of at least three times daily in the past 30 days among adolescents in WP Labuan, 2022

**Table 3.3.4** Prevalence of fruits and vegetables intake of at least five times daily in the past 30 days among adolescents in WP Labuan, 2022

**Table 3.3.5** Prevalence of adolescents who did not consume fruit, vegetable or both in the past 30 days in WP Labuan, 2022

**Table 3.3.6** Prevalence of carbonated soft drinks intake of at least once a day in the past 30 days among adolescents in WP Labuan, 2022

**Table 3.3.7** Prevalence of plain water intake of less than 6 glasses daily in the past 30 days among adolescents in WP Labuan, 2022

**Table 3.3.8** Prevalence of milk and milk products intake of at least twice daily in the past 30 days among adolescents in WP Labuan, 2022

**Table 3.3.9** Prevalence of fast-food intake of at least 3 days in the past 7 days among adolescents in WP Labuan, 2022

### NUTRITIONAL STATUS

**Table 3.4.1** Prevalence of stunting (HAZ <-2SD) among adolescents in WP Labuan, 2022

**Table 3.4.2** Prevalence of thinness (BAZ <-2SD) among adolescents in WP Labuan, 2022

**Table 3.4.3** Prevalence of overweight (BAZ >+1SD to ≤+2SD) and obesity (BAZ >+2SD) among adolescents in WP Labuan, 2022

### DRUG USE

**Table 3.5.1** Prevalence of ever drug use among adolescents in WP Labuan, 2022

**Table 3.5.2** Prevalence of current drug use among adolescents in WP Labuan, 2022

**Table 3.5.3** Prevalence of inhalant use among adolescents in WP Labuan, 2022

**Table 3.5.4** Prevalence of first use of drug before the age of 14 years among ever used drug, adolescents in WP Labuan, 2022

### ORAL AND HAND HYGIENE

**Table 3.6.1** Prevalence of poor or very poor perception of oral health among adolescents in WP Labuan, 2022

**Table 3.6.2** Prevalence of teeth brushing 2 times a day in the past 30 days among adolescents in WP Labuan, 2022

**Table 3.6.3** Prevalence of never did tongue cleaning among adolescents in WP Labuan, 2022

**Table 3.6.4** Prevalence of did not know if their toothpaste is fluoridated among adolescents in WP Labuan, 2022

**Table 3.6.5** Prevalence of use of dental floss among adolescents in WP Labuan, 2022

**Table 3.6.6** Prevalence of last dental visit in the past 12 months among adolescents in WP Labuan, 2022

**Table 3.6.7** Prevalence of having missed classes or online learning among adolescents with toothache in WP Labuan, 2022

LIST OF TABLES

**Table 3.6.8** Prevalence of avoidance of smiling due to teeth appearance among adolescents in WP Labuan, 2022

**Table 3.6.9** Prevalence of using soap most of the time or always during handwashing in the past 30 days among adolescents in WP Labuan, 2022

**Table 3.6.10** Prevalence of handwashing most of the time or always before eating in the past 30 days among adolescents in WP Labuan, 2022

**Table 3.6.11** Prevalence of handwashing most of the time or always after using the toilet in the past 30 days among adolescents in WP Labuan, 2022

**Table 3.6.12** Prevalence of handwashing using running water before eating at school in the past 30 days among adolescents in WP Labuan, 2022

**MENTAL HEALTH PROBLEMS**

**Table 3.7.1** Prevalence of loneliness “most of the time or always” in the past 12 months among adolescents in WP Labuan, 2022

**Table 3.7.2** Prevalence of inability to sleep “most of the time or always” due to worry in the past 12 months among adolescents in WP Labuan, 2022

**Table 3.7.3** Prevalence of suicidal ideation in the past 12 months among adolescents in WP Labuan, 2022

**Table 3.7.4** Prevalence of suicidal plan in the past 12 months among adolescents in WP Labuan, 2022

**Table 3.7.5** Prevalence of suicidal attempt “at least once” in the past 12 months among adolescents in WP Labuan, 2022

**Table 3.7.6** Prevalence of not having any close friend among adolescents in WP Labuan, 2022

**Table 3.7.7** Prevalence of depression among adolescents in WP Labuan, 2022

**PHYSICAL ACTIVITY**

**Table 3.8.1** Prevalence of being physically active (at least 60 minutes daily) for a total of 5 days or more in the past 7 days among adolescents in WP Labuan, 2022

**Table 3.8.2** Prevalence of active commuting to school (walk or ride a bicycle to or from school for at least 3 days or more in the past 7 days) among adolescents in WP Labuan, 2022

**Table 3.8.3** Prevalence of spending at least 3 hours in sitting activities among adolescents in WP Labuan, 2022

**PROTECTIVE FACTORS**

**Table 3.9.1** Prevalence of truancy in the past 30 days among adolescents in WP Labuan, 2022

**Table 3.9.2** Prevalence of having peer support in the past 30 days among adolescents in WP Labuan, 2022

**Table 3.9.3** Prevalence of having parental or guardian supervision in the past 30 days among adolescents in WP Labuan, 2022

**Table 3.9.4** Prevalence of having parental or guardian connectedness in the past 30 days among adolescents in WP Labuan, 2022

**Table 3.9.5** Prevalence of having parental or guardian bonding in the past 30 days among adolescents in WP Labuan, 2022

**Table 3.9.6** Prevalence of having parental or guardian respect for privacy in the past 30 days among adolescents in WP Labuan, 2022

**SEXUAL BEHAVIOURS**

**Table 3.10.1** Prevalence of ever had sexual intercourse among adolescents in WP Labuan, 2022

**Table 3.10.2** Prevalence of current sexual intercourse in the past 30 days among adolescents in WP Labuan, 2022

**Table 3.10.3** Proportion of sexual practices among those who ever had sex among adolescents in WP Labuan, 2022

**Table 3.10.4** Prevalence of adequate HIV knowledge among adolescents by Sociodemographic Characteristics in WP Labuan, 2022

**Table 3.10.5** Percentage of correct responses by item of UNGASS indicator among adolescents in WP Labuan, 2022

**TOBACCO USE**

**Table 3.11.1** Prevalence of current any tobacco uses among adolescents in WP Labuan, 2022

**Table 3.11.2** Prevalence of e-cig/vape use among adolescents in WP Labuan, 2022

**Table 3.11.3** Proportion of source of cigarette obtaining during the last time smoking in the past 30 days among adolescents in WP Labuan, 2022

**Table 3.11.4** Proportion of source of e-cig/vape obtaining during the last time using e-cig/vape in the past 30 days among adolescents in WP Labuan, 2022

**Table 3.11.5** Prevalence of exposure to second-hand smoke among adolescents in WP Labuan, 2022

**Table 3.11.6** Prevalence of currently seeing or noticing any tobacco product advertising or promotion at the point of sale in the past 30 days among adolescents in WP Labuan, 2022

## LIST OF TABLES

## VIOLENCE AND UNINTENTIONAL INJURY

<b>Table 3.12.1</b>	Prevalence of involvement in violence at least once in the past 12 months among adolescents in WP Labuan, 2022
<b>Table 3.12.2</b>	Prevalence of had serious injury at least once in the past 12 months among adolescents in WP Labuan, 2022
<b>Table 3.12.3</b>	Major cause of the most serious injury sustained in the past 12 months among those who were injured, adolescents in WP Labuan, 2022
<b>Table 3.12.4</b>	Prevalence of being abused at least once in the past 30 days among adolescents in WP Labuan, 2022
<b>Table 3.12.5</b>	Prevalence of experience in being bullied at least once in the past 30 days among adolescents in WP Labuan, 2022
<b>Table 3.12.6</b>	Most common ways of being bullied at least once in the past 30 days among adolescents in WP Labuan, 2022
<b>Table 3.12.7</b>	Prevalence of involvement in cyberbullying activities (perpetrator) a few times within a year or more among adolescents in WP Labuan, 2022
<b>Table 3.12.8</b>	Most common ways of involvement in cyberbullying activities (perpetrator) a few times within a year or more among adolescents in WP Labuan, 2022

## IMPACT OF COVID-19

<b>Table 3.13.1</b>	Adolescents' perspectives on the impact of the COVID-19 pandemic on their family: Self-reported findings among adolescents in WP Labuan, 2022
---------------------	---


**LIST OF APPENDICES**


---

Appendix 1	Members of Steering Committee NHMS 2019-2022
Appendix 2	Terms of Reference for NHMS 2022 Steering Committee
Appendix 3	List of members of Central Coordinating Committee, NHMS 2022
Appendix 4	Terms of Reference for NHMS 2022 Central Coordinating Team (CCT)
Appendix 5	List of Research Team Members, NHMS 2022
Appendix 6	List of Data Collection Teams, NHMS 2022

## EXECUTIVE SUMMARY

The Adolescent Health Survey (AHS) 2022 was conducted from June to July 2022 with the aim of determining the prevalence of health risk behaviours and protective factors among adolescents in Malaysia. This nationwide cross-sectional survey used a two-stage stratified sampling design and a validated self-administered questionnaire. Out of 2798 secondary schools under the Ministry of Education (MOE) and the Ministry of Rural and Regional Development (MARA), 240 schools were randomly selected and a total of 37,479 students were eligible to participate in the survey. The findings showed that a total of 239 schools with 33,523 adolescents were involved in this study, resulting in an overall response rate of 89.0%. In WP Labuan, 8 secondary schools were randomly selected, and out of 2267 eligible students, 2033 students completed the survey, yielding a response rate of 89.7%.

### WP Labuan Key Findings

The study revealed that the prevalence of current use of any tobacco products, and current e-cig/vape users among adolescents in WP Labuan was 22.2% and 19.2% respectively. The prevalence of ever-alcohol drinkers among adolescents in WP Labuan was 19.3%, 64.1% of them had their first alcoholic beverage before the age of 14 years. The prevalence of ever having sex and had sex in the past 30 days among adolescents was 7.1% and 4.5%, respectively. Of those who ever had sex, 39.1% had their first sexual experience before age 14, and 8.5% had at least two sexual partners. Only 9.1% of respondents or their partners had used condoms, while 12.7% used other birth control methods. A total of 20.2% of adolescents had been seriously injured in the past 12 months, with the two most common causes of injury being falls and motor vehicle accidents. Among respondents, 17.4% claimed to have been physically attacked in the past 12 months, while 20.5% of adolescents claimed to have been involved in physical fights. With regards to bullying, 9.9% reported having been bullied in the past 30 days. A total of 20.3% of adolescents in WP Labuan reported feeling lonely, and 17.9% reported being unable to sleep "most of the time or always" due to worry in the 12 months prior to the survey. Prevalence of suicidal ideation, plan and attempt were 17.7%, 14.0%, and 13.0%, respectively. Overall, 34.4% of adolescents reported being depressed. The prevalence of truancy among adolescents in the past 30 days was 29.8%, and only 47.5% claimed to have peer support. Adolescents who reported having parental or guardian supervision, parental or guardian connectedness and parental or guardian bonding were 8.1%, 18.9% and 27.2%, respectively. Overall, 85.7% of adolescents reported brushing their teeth twice a day in the past 30 days. A total of 43.3% of adolescents reported not knowing whether their toothpaste contained fluoride while only 24.1%

used dental floss. In the past 30 days, 72.8% always used soap when washing their hands, 84.9% always washed their hands before eating, and 87.5% reported that they always washed their hands after using the toilet. In relation to dietary behaviours, 4.1% reported being hungry most of the time or always in the past 30 days because there was not enough food at home. The consumption of fruits at least twice daily was 35.5% and vegetables at least thrice daily was 25.6% in the past 30 days. Consumption of carbonated drinks at least once daily in the past 30 days was reported at 46.5%, while 15.1% consumed food from fast food restaurants for at least three days in the past seven days. The prevalence of stunting and thinness among adolescents was 7.8% and 6.1%, respectively, while the prevalence of overweight was 17.11% and obesity was 15.9%. Prevalence of being physically active was 22.4% and 17.8% of adolescents reported active transportation to school. In addition, 66.7% of adolescents had spent at least three hours on a typical or usual day in sitting activities. Overall, 3.9% reported had ever used drug and the prevalence of current drug users was 2.1% while 1.0% reported had ever used kratom.

### Malaysia Key Findings

The study revealed that the prevalence of current use of any tobacco products, current tobacco smokers, current cigarette smokers and current e-cig/vape users among adolescents in Malaysia was 18.5%, 9.0%, 6.2% and 14.9% respectively. Among those who smoked cigarettes and among those who used e-cig/vape, 65.7% had initiated cigarette smoking, and 48.5% had initiated e-cig/vape use, respectively, before the age of 14 years. The prevalence of current alcohol drinkers among adolescents was 7.4%. While the prevalence of ever-alcohol drinkers among adolescents in Malaysia was 18.6%, 64.6% of them had their first alcoholic beverage before the age of 14 years.

The prevalence of ever having sex and had sex in the past 30 days among adolescents was 7.6% and 5.7%, respectively. Of those who ever had sex, 32.8% had their first sexual experience before age 14, and 10.7% had at least two sexual partners. Only 11.8% of respondents or their partners had used condoms, while 11.9% used other birth control methods. A total of 20.4% of adolescents had been seriously injured in the past 12 months, with the two most common causes of injury being falls and motor vehicle accidents. Among respondents, 14.8% claimed to have been physically attacked in the past 12 months, while 16.0% of adolescents claimed to have been involved in physical fights. With regards to bullying, 8.6% reported having been bullied in the past 30 days.

A total of 16.2% of adolescents in Malaysia reported feeling lonely, and 4.2% said that they had no close friends. A total of 12.9% reported being unable to sleep “most of the time or always” due to worry in the 12 months prior to the survey. Prevalence of suicidal ideation, plan and attempt were 13.1%, 10.0%, and 9.5%, respectively. Overall, 26.9% of adolescents reported being depressed. The prevalence of truancy among adolescents in the past 30 days was 25.6%, and only 46.0% claimed to have peer support. Adolescents who reported having parental or guardian supervision, parental or guardian connectedness and parental or guardian bonding were 9.9%, 24.2% and 33.4%, respectively. Overall, 82.2% of adolescents reported brushing their teeth twice a day in the past 30 days. A total of 43.3% of adolescents reported not knowing whether their toothpaste contained fluoride while only 21.4% used dental floss. In the past 30 days, 69.3% always used soap when washing their hands, 84.5% always washed their hands before eating, and 86.5% reported that they always washed their hands after using the toilet.

In relation to dietary behaviours, 2.5% reported being hungry most of the time or always in the past 30 days because there was not enough food at home. The consumption of fruits at least twice daily was 37.3% and vegetables at least thrice daily was 27.1% in the past 30 days. Consumption of carbonated drinks at least once daily in the past 30 days was reported at 32.4%, while 10.6% consumed food from fast food restaurants for at least three days in the past seven days. The prevalence of stunting and thinness among adolescents was 6.8% and 8.3%, respectively, while the prevalence of overweight was 16.2% and obesity was 14.3%. Prevalence of being physically active for a total of at least 60 minutes daily for five days or more in the past seven days was 21.4% and 27.0% of adolescents reported active transportation to school. In addition, 66.7% of adolescents had spent at least three hours on a typical or usual day in sitting activities.

### Recommendations:

In view of the above findings, the following recommendations are suggested:

- Strengthening the multi-approach school-based nutrition and physical activity intervention to motivate behaviour modification for improving healthy eating and lifestyle amongst adolescents.
- Improving the national school curriculum that teaches life skills such as effective coping strategies as part of “Program Minda Sihat”.
- A more comprehensive sexual and reproductive health education programmes should be planned and executed among adolescents.
- Strengthening the current law and taking legal action in controlling the accessibility of tobacco products.

## 1.0 INTRODUCTION

Adolescence is a life phase in which the opportunities for health are great and future patterns of adult health are established; it is a critical stage in life with significant physical, emotional, cognitive, and social development and other disruptions in their communities.<sup>1</sup> As much as one-third of the global Burden of Disease (BOD) is attributable to adolescent behavioural choices and events.<sup>2</sup> In order to improve adolescent health globally, the World Health Organization (WHO) has initiated the development of the health risk behaviours measurement tools known as the Global School-based Student Health Survey (GSHS).<sup>3</sup> More than 140 countries have used the GSHS to periodically monitor the prevalence of important health risk behaviours and protective factors among adolescents.<sup>4</sup> In Malaysia, adolescents comprise approximately 15.6% of the total Malaysian population, and the national data on health risks and behaviours are fundamental in developing policies and programmes for adolescents. Thus, the Ministry of Health, Malaysia took a step forward in collaborating with the WHO to conduct the first GSHS Malaysia in 2012 among adolescents aged 13 to 17 years, which aimed to determine the baseline of the health status of adolescents in Malaysia.<sup>5</sup> In 2017, the second adolescent health study (AHS) using the GSHS methods and questionnaire was conducted in the country.<sup>6</sup> These surveys revealed an increasing trend of health risk behaviours among adolescents in Malaysia.<sup>5,6</sup> With the increasing trend of non-communicable disease risk factors and other behaviour-related risks, it is timely for the survey to be repeated in 2022 to further monitor the health status of adolescents in the country. The Ministry of Health conducted this third national survey on adolescents with the co-operation from the Ministry of Education to determine the prevalence of health risk behaviours and protective factors among adolescents in Malaysia.

### 1.1 Objectives

#### 1.1.1 General Objectives

To determine the prevalence of health risk behaviours and protective factors among adolescents in Malaysia.

#### 1.1.2 Specific Objectives

To determine the prevalence of:

- i. Alcohol use
- ii. Dietary behaviours
- iii. Drug use
- iv. Hygiene (including oral health)
- v. Mental health problems
- vi. Physical activity
- vii. Protective factors
- viii. Sexual behaviours
- ix. Tobacco use
- x. Violence and unintentional injury
- xi. Adolescents' perspectives on the impact of the COVID-19 pandemic on their families

## 2.0 METHODOLOGY

### 2.1 Study Design

The National School-Based Student Health Survey 2022 was a nationwide cross-sectional study of secondary school adolescents in Malaysia.

### 2.2 Sampling Frame and Target Population

The sampling frame comprised national secondary schools registered in 2021, which include government schools and private schools under the purview of the Ministry of Education (MOE) and the Ministry of Rural and Regional Development (MARA). According to the frame, there were 2798 secondary schools in Malaysia (**Table 2.1**). An equal proportion was sampled from 13 States and three Federal Territories to represent adolescents in each State / Federal Territories. The target population was secondary school adolescents aged between 13 to 17 years studying in form 1 until form 5 based on the local school categorization.

### 2.3 Sample Size Calculation

The sample size was calculated based on the objectives of each module using the sample size calculation formula for a single proportion. The sample size calculation was based on a few criteria, as stated below:

$$n_0 = \frac{z_{\alpha/2} p(1-p)}{e^2}$$

Where:

- Variance of proportion of the variable of interest (Based on AHS 2017 survey)
- Margin of error (e) (Between 0.01 to 0.05)
- Confidence interval of 95%

To ensure optimum sample size to estimate the prevalence of the health conditions specified in the survey with acceptable precision, a few adjustments were made to the sample size calculation as follows:

- design effect (deff) of 2,
- nonresponse rates of 20%, and
- The sample size was then adjusted according to the need of the analysis, whether the estimates were going to be done at the national or the state level.

Thus, the final sample sizes for adolescents at national and state levels were 36,000 and 2250, respectively (**Table 2.1**)

**Table 2.1: Distribution of secondary schools sampled, by state**

No.	State / Federal Territories	Total Number of Schools	Number of Schools Sampled	Number of adolescents sampled
1	Johor	328	16	2250
2	Kedah	219	16	2250
3	Kelantan	189	16	2250
4	Melaka	88	16	2250
5	N. Sembilan	142	16	2250
6	Pahang	211	16	2250
7	Pulau Pinang	148	16	2250
8	Perak	276	16	2250
9	Perlis	33	16	2250
10	Selangor	380	16	2250
11	Terengganu	166	16	2250
12	Sabah	245	16	2250
13	Sarawak	214	16	2250
14	WP Kuala Lumpur	135	16	2250
15	WP Labuan	12	8	2250
16	WP Putrajaya	12	8	2250
<b>Total</b>		<b>2798</b>	<b>240</b>	<b>36000</b>

### 2.4 Sampling Design

The country was stratified according to the 16 states, including federal territories, for the sampling. A multistage stratified cluster sampling method was used, and it involved two stages. The first stage was the selection of secondary schools from all eligible schools in Malaysia. Subsequently, the 240 schools were selected randomly with probability proportional to enrolment (PPS) in forms 1, 2, 3, 4, and 5. In each state, 16 secondary schools were selected, except for 2 smaller federal territories (Labuan, Putrajaya - 8 schools each) (**Table 2.1**). The second stage involved the selection of classes (secondary sampling units). All classes in forms 1, 2, 3, 4, and 5 were included in the sampling frame. Systematic probability sampling with a random start was used to select classes from each selected school. All adolescents in the selected classes were invited to involve in the survey.

### 2.5 Ethical Approval and Consent Forms

Ethical approval was obtained from the Medical and Research Ethics Committee (MREC), Ministry of Health, Malaysian (NMRR-21-157-58261). The permission to conduct the study was obtained from the Ministry of Education at the national, state and school levels. Only consented adolescents with consented parents were included in the study. Their participation in the study was voluntary.



## 2.6 Study Instrument

A validated self-administered questionnaire was used for data collection in NHMS 2022. The questionnaires were translated into the Malay, Chinese and Tamil languages and back-translated to English to ensure the quality of the translation. The questionnaires were then field-tested, revised, finalised, and approved by the NHMS 2022 Questionnaire Review Committee. The questionnaire consisted of 10 core modules and 1 additional module, which included the following topics:

- Alcohol use
- Dietary behaviours
- Drug use
- Hygiene (including oral health)
- Mental health problems
- Physical activity
- Protective factors
- Sexual behaviours
- Tobacco use
- Violence and unintentional injury
- Adolescents' perspectives on the impact of the COVID-19 pandemic on their families

## 2.7 Data Collection

This cross-sectional survey was conducted from June to July 2022 among adolescents in forms 1, 2, 3, 4, and 5 across Malaysia by 34 data collection teams: two teams for each state in Peninsular Malaysia, including Labuan and three teams for Sabah and Sarawak. Each state was assigned a field supervisor to oversee survey activities. A one-week training workshop was conducted for the field supervisor and 133 temporary data collectors before data collection. After completing the training, the assigned field supervisor and data collection teams travelled to their respective sites to conduct the survey at the selected schools. The adolescents answered the questionnaires on the optical mark recognition (OMR) answer sheet. The team leader verified the OMR sheets before posting them to Institute for Public Health (IPH).

## 2.8 Quality Control

Quality control of the whole survey was done at various stages. During the planning stage, quality was ensured through a robust survey design, validated questionnaires and tools, manuals, and standardised training. In the field, the team leader and field supervisor checked the quality of the data collected. At the same time, members of the Central Coordinating Team (CCT) at IPH monitored data collection progress and conducted data quality control on a weekly basis. Figure 1 detailed the organization chart at Institute for Public Health level.

## 2.9 Data Processing and Quality Centre

All data processing and quality activities were centralised at IPH, starting from receiving the OMR bundles from the field until the handover of the dataset to the data analysis team. Four stations were set up at this Centre to ensure the activity ran smoothly (Figure 2).

## 2.10 Data Analysis

SPSS version 26.0 was used for data analysis. The data was examined for quality control and cleaned for any inconsistencies. Analysis was done according to objectives, working definition and dummy tables prepared by each research team. A complex sample analysis procedure was performed with a 95% confidence interval. Prevalence and percentages were used to illustrate the findings of each scope.

## 2.11 References

1. Sawyer SM, Afifi RA, Bearinger LH, et al. Adolescence: A foundation for future health. *Lancet* 2012;379:1630e40
2. Guthold R, Moller AB, Azzopardi P, Ba MG, Fagan L, Baltag V, Say L, Banerjee A, Diaz T. The Global Action for Measurement of Adolescent health (GAMA) Initiative-Rethinking Adolescent Metrics. *J Adolesc Health*. 2019 Jun;64(6):697-699
3. World Health Organization (WHO). WHO, Global school-based student health survey (GSHS). WHO. <http://www.who.int/chp/gshs/en/>. Accessed 17 Aug 2022
4. Biswas T, Townsend N, Huda M, Maravilla J, Begum T, Pervin S, et al. 2022. Prevalence of multiple non-communicable diseases risk factors among adolescents in 140 countries: A population-based study. *eClinicalMedicine*. 2022;52: 101591
5. Institute for Public Health (IPH) 2012. National Health and Morbidity Survey (NHMS) 2012: Adolescent Health Survey 2012, Malaysia
6. Institute for Public Health (IPH) 2017. National Health and Morbidity Survey (NHMS) 2017: Adolescent Health Survey 2017, Malaysia




Figure 1: Organisation chart for data collection team NHMS 2022




Figure 2: Workflow of Data Processing NHMS 2022

## 3.0 FINDINGS

### 3.1 General Findings

Overall, 239 out of 240 schools participated in the survey, resulting in a 99.6% school response rate. The response rate for classes was 100% and the student's response rate was 89.4% (n=33,523). Thus, the overall response rate was 89.0%. Based on the state, the highest number of adolescents who participated in the survey were from Terengganu (95.9%), and the lowest was WP Putrajaya (83.4%) (Table 3.1.1). The geographic information system (GIS) on the mapping of selected secondary schools is shown in Figure 3.

Table 3.1.1: Response rate at student level, by state, 2022

State	Selected Schools	Eligible Adolescents	Completed OMR forms	Response Rate (%)
Johor	16	2336	2005	85.83
Kedah	16	2312	2172	93.94
Kelantan	16	2368	2138	90.29
Melaka	16	2373	1986	83.69
N. Sembilan	16	2422	2210	91.25
Pahang	16	2382	2171	91.14
Pulau Pinang	16	2300	2044	88.87
Perak	16	2384	2126	89.18
Perlis	16	2160	2004	92.78
Selangor	16	2366	2048	86.56
Terengganu	16	2314	2219	95.89
Sabah	16	2342	2086	89.07
Sarawak	16	2442	2189	89.64
WP Kuala Lumpur	16	2338	2114	90.42
WP Labuan	8	2267	2033	89.68
WP Putrajaya	8	2373	1978	83.35
<b>Total</b>	<b>240</b>	<b>37479</b>	<b>33523</b>	<b>89.44</b>




Figure 3: GIS mapping of the selected secondary schools

## 3.2 Alcohol Consumption

Contributors: Tania Gayle Robert, Hamizatul Akmal Abd Hamid, Mohd Hatta Abdul Mutalip, Chong Zhuo Lin, Halizah Mat Rifin, Norli Abdul Jabbar, Rusdi Abd Rashid, Thamil Arasu Saminathan, Muhammad Fadhli Mohd Yusoff, Faizul Akmal Abdul Rahim, Mohd Ruhaizie Riyadzi

### 3.2.1 Introduction

Annually, the harmful use of alcohol results in approximately 3 million deaths worldwide<sup>1</sup>. Alcohol is responsible for 5.1% of the global burden of disease and injury, as measured by disability-adjusted life years (DALYs)<sup>1</sup>. It is the main cause of premature death and disability in individuals aged 15 to 49, accounting for 10% of all deaths in this age group<sup>2</sup>.

### 3.2.2 Objectives

- i. To determine the prevalence of ever and current drinkers among adolescents
- ii. To describe the socio-demographic characteristics of ever and current drinkers among adolescents
- iii. To identify the age of alcohol drinking initiation among adolescents
- iv. To identify the sources of obtaining alcoholic beverages among adolescents
- v. To identify the prevalence of drunkenness among adolescents who consume alcohol
- vi. To determine the frequency of social problems related to alcohol consumption among adolescents

### 3.2.3 Variable definitions

- **Drinking alcohol:** A “drink” is a glass of wine, tuak, lihing, bahar, ijuk or toddy; a can of beer, a small glass of liquor’ or mixed drink. Drinking alcohol does not include drinking a few sips of wine for religious purposes.
- **Ever drinkers:** Those who had a history of alcohol consumption in their lifetime.
- **Current drinkers:** Those who had at least a “drink” of alcohol in the past 30 days.
- **Drunkenness:** When someone demonstrates signs such as staggering when walking, not being able to speak right and throwing up after consuming alcohol in a lifetime.
- **Social problems:** Having trouble with family or friends, missed school or got into fights as a result of drinking alcohol in a lifetime.

### 3.2.4 Findings

#### Ever Alcohol Drinkers

The prevalence of ever alcohol drinkers among adolescents in WP Labuan was 19.3% (95% CI: 12.85, 27.95). Males had a higher prevalence of 20.3% (95% CI: 11.78, 32.69) compared to females of 18.3% (95% CI: 11.15, 28.67). (Table 3.2.1)

#### Initiation of First Alcohol Use Before 14 years old

Among ever alcohol drinkers, 64.1% had their first alcoholic beverage before the age of 14 years. (Table 3.2.3)

#### Sources of Obtaining Alcoholic Beverages Among Current Alcohol Drinkers

Among current alcohol drinkers, 56.3% of adolescents obtained their alcoholic beverages from their family, followed by 13.9% of them got it from friend. (Table 3.2.4)

#### Social Problems as a Result of Alcohol Drinking

Overall, the majority (81.8%) of the adolescents did not get into trouble with family or friends, missed school or got into a fight as a result of drinking alcohol among ever alcohol drinkers. (Table 3.2.5)

#### Parental and peer drinking

Parental drinking among current alcohol drinkers in WP Labuan was 63.2%, while peer drinking was 84.3%.

#### Drunkenness

Among the ever drinkers, 35.5% reported drunkenness.

### 3.2.5 Discussion / Conclusion

According to this study, the prevalence of ever alcohol drinkers among adolescents in WP Labuan in 2022 is 19.3%. However, at 64.1%, the proportion of adolescents who drank alcohol for the first time before turning 14 years old remained high. As in surveys, family members remained the primary source of acquiring alcoholic beverages. Majority of the adolescents did not get into trouble with family or friends, missed school or got into a fight as a result of drinking alcohol. The percentage of peer drinking among the current drinkers is the most contribution of adolescent involvement in alcohol use in WP Labuan, which stood at 84.3%.

### 3.2.6 Recommendations

Given that adolescence is the age when adult habits and social standards are formed, it is essential to look into the issue of adolescent drinking. The detrimental effects of alcohol drinking at a young age should be made clear to parents and other caregivers through mass media. Enhance school-based prevention programmes in school with high rates of alcohol consumption to enable teachers to assess pupils for alcohol consumption and launch early intervention for such adolescents.

### 3.2.7 References

1. World Health Organization, Key Fact: Alcohol. <https://www.who.int/news-room/fact-sheets/detail/alcohol>
2. World Health Organisation. Alcohol. <https://www.who.int/health-topics/alcohol>

**Table 3.2.1: Prevalence of ever alcohol drinkers among adolescents in WP Labuan, 2022**

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
<b>WP LABUAN</b>	395	1254	19.3	12.85	27.95
<b>Sex</b>					
Male	207	643	20.3	11.78	32.69
Female	188	611	18.3	11.15	28.67
<b>Form</b>					
Form 1	66	231	15.8	8.89	26.53
Form 2	76	262	17.6	9.78	29.52
Form 3	73	257	19.4	12.71	28.42
Form 4	89	256	21.9	11.34	38.06
Form 5	91	248	23.6	12.63	39.78
<b>Ethnicity</b>					
Malay	62	194	6.7	4.60	9.53
Chinese	100	329	42.3	32.03	53.23
Indian	1	-	-	-	-
Bumiputera Sabah	164	515	28.4	16.81	43.71
Bumiputera Sarawak	25	77	31.0	18.53	47.00
Others	43	135	18.6	10.97	29.89

- Prevalence with high RSE, not reported

**Table 3.2.2: Proportion of ever alcohol drinkers according to alcohol initiation age among adolescents in WP Labuan, 2022**

Initiation age of alcohol drinking	Unweighted count	Percentage (%)
Below 14 years old	205	64.1
14 years old and above	126	35.9

**Table 3.2.3: Usual sources of obtaining alcohol in the past 30 days among current drinkers among adolescents in WP Labuan, 2022**

Sources of obtaining alcohol	Unweighted count	Percentage (%)
I bought from a store, shop or from a street vendor	29	-
I gave someone else money to buy it for me	3	-
I got it from my friend	20	13.9
I got it from my family	71	56.3
I stole it or got it without permission	0	-
I got it some other way	8	-

- Prevalence with high RSE, not reported

**Table 3.2.4: Number of times (got into trouble with family or friends, missed school or got into a fight as a result of drinking alcohol) among ever alcohol drinkers (proportion) among adolescents in WP Labuan, 2022**

Number of times	Unweighted count	Percentage (%)
0 times	323	81.8
1 to 2 times	49	12.2
3 to 9 times	13	3.4
10 or more times	10	-

- Prevalence with high RSE, not reported

### 3.3 Dietary Behaviours

Contributors: Lai Wai Kent, Suhaila Abdul Ghaffar, Chong Chean Tat, Munawara Pardi, Nurul Huda Ibrahim, Syafinaz Mohd Sallehuddin, Siti Adibah Ab. Halim, Ainan Nasrina Ismail, Teh Wai Siew

#### 3.3.1 Introduction

The changes in dietary practices are required across all age ranges, but adolescence should be a focus of particular attention because the changes in lifestyle and the development of dietary habits during that stage of life have striking effects<sup>1</sup>. Poor dietary intake during this life stage is closely related to overweight and obesity, and unhealthy eating practices which lead to detrimental health effects later in life<sup>2</sup>. Therefore, a study was conducted to examine dietary practices with regards to fruit and vegetable intakes, carbonated drinks, plain water, milk and milk products intake and fast-food consumption among adolescents in WP Labuan. Establishing and consuming a nutrient-dense diet during the transition from adolescence into young adulthood may protect against future chronic diseases, promote optimal health outcomes and to prevent excess weight gain<sup>3</sup>.

#### 3.3.2 Objectives

- i. To describe the prevalence of adolescents who had gone hungry in the past 30 days
- ii. To describe the prevalence of fruit intakes of at least twice daily in the past 30 days among adolescents
- iii. To identify the prevalence of vegetable consumption of at least three times daily in the past 30 days
- iv. To identify the prevalence of fruit and vegetables consumption of at least five times daily in the past 30 days
- v. To describe the prevalence of carbonated drink intake of at least once a day in the past 30 days
- vi. To describe the prevalence of plain water intake of less than 6 glasses per day in the past 30 days
- vii. To identify the prevalence of milk and milk product intakes of at least two times daily in the past 30 days
- viii. To identify the prevalence of fast-food consumption of at least three days in the past 7 days

#### 3.3.3 Variable definitions

- **Gone Hungry:** Adolescents who had gone hungry most of the time or always because there was not enough food at home for the past 30 days, or living without financial means to access enough food for active and healthy living.
- **Fruit intakes:** Fruits intake of at least twice daily in the past 30 days, inclusive all types of fruits.
- **Vegetable intakes:** Vegetable intakes of at least three times daily in the past 30 days.
- **Plain water intake:** Includes mineral water, boiled water or tap water

- **Carbonated drinks intake:** carbonated drinks consumption of at least once daily in the past 30 days.
- **Dairy product intake:** milk and milk product intakes at least two times daily in the past 30 days.
- **Fast food intake:** Consuming food from fast food outlets at least three days in the past seven days.

#### 3.3.4 Findings

##### Gone hungry

About 4.1% (95% CI: 2.97, 5.66) of adolescents reported being hungry most of the time or always because there was not enough food at home in the past 30 days (**Table 3.3.1**). Prevalence of being hungry was higher among females (5.2%, 95% CI: 4.00, 6.85) as compared to male counterparts (2.9%, 95% CI: 4.00, 6.85).

##### Fruit consumption

A total of 35.5% (95% CI: 34.12, 36.89) of adolescents consumed fruit at least twice daily in the past 30 days (**Table 3.3.2**). Males (36.9%, 95% CI: 33.25, 40.62) reported higher fruit intake than females (34.2%, 95% CI: 32.22, 36.23).

##### Vegetable consumption

About 25.6% (95% CI: 23.76, 27.59) of adolescents consumed vegetables at least three times daily in the past 30 days (**Table 3.3.3**). Males reported significantly higher vegetable intake (28.1%, 95% CI: 26.11, 30.21) compared to females (23.3%, 95% CI: 21.02, 25.58).

##### Fruits and vegetables intake

About 15.9% (95% CI: 14.19, 17.72) of adolescents consumed fruits and vegetables at least five times daily in the past 30 days (**Table 3.3.4**). Males (17.1%, 95% CI: 14.17, 20.46) reported higher consumption of fruits and vegetables than females (14.7%, 95% CI: 13.62, 15.89).

##### Never Consume fruit

About 13.4% (95% CI: 8.11, 16.09) of adolescents reported never consume fruit in the past 30 days (**Table 3.3.5**). Prevalence of never consume fruit was 12.6% (95% CI: 9.81, 16.09) among males and 14.2% (95% CI: 11.36, 17.65) among females.

##### Never Consume Vegetable

About 11.7% (95% CI: 9.11, 14.93) of adolescents never consume vegetable in the past 30 days (**Table 3.3.5**). Prevalence of never consume vegetable was 12.1% (95% CI: 8.43, 16.98) among males and 11.4% (95% CI: 8.78, 14.62) among females.

##### Never Consume Fruit and Vegetable

A total of 3.3% (95% CI: 2.50, 4.31) of adolescents never consume fruit and vegetable in the past 30 days (**Table 3.3.5**). Prevalence of never consume fruit and vegetable was 3.8% (95% CI: 2.47, 5.87) among males and 2.8% (95% CI: 2.09, 3.69) among females.



**Carbonated soft drinks intake**

Overall, 46.5% (95% CI: 42.12, 50.89) of adolescents consumed carbonated soft drinks at least once daily in the past 30 days (Table 3.3.6). Males (46.6%, 95% CI: 40.27, 53.03) consumed carbonated soft drinks more frequently than females (46.4%, 95% CI: 41.23, 51.59).

**Plain water intake**

About 44.9% (95% CI: 42.76, 47.04) adolescents drank plain water less than six glasses per day in the past 30 days (Table 3.3.7). Prevalence of plain water intake of less than six glasses per day among females (51.4%, 95% CI: 48.18, 54.65) was significantly higher than males (38.0%, 95% CI: 34.83, 41.31).

**Milk and milk products intake**

About 25.7% (95% CI: 22.97, 28.56) of adolescents consumed milk/milk products at least two times per day in the past 30 days (Table 3.3.8). Findings showed that the prevalence of milk/milk products consumption among females (25.6%, 95% CI: 21.83, 31.93) was slightly higher compared to males (24.7%, 95% CI: 21.07, 28.75)

**Fast food intake**

About 15.1% (95% CI: 13.30, 17.12) of adolescents consumed fast food at least three days in the past seven days (Table 3.3.9). The prevalence of fast-food intake was significantly higher among females (17.9%, 95% CI: 15.68, 20.38) compared to males (12.2%, 95% CI: 9.69, 15.17).

**3.3.5 Discussions / Conclusion**

There were 4.1% of adolescents who reported being hungry due to lack of food at homes. Prevalence of adolescents reported consuming fruits at least twice daily and vegetables at least three times daily was 35.5% and 25.6%, respectively. However, only 15.9% of them consumed fruits and vegetables five times daily. About 44.9% of adolescents reported drinking plain water less than 6 glasses and 25.7% consumed milk/milk products at least two times daily. The consumption of carbonated soft drinks of at least once daily in the past 30 days was reported at 46.5% while 15.1% consumed food from fast food restaurants for at least three days in the past seven days.

**3.3.6 Recommendations**

The research finding shows that there is a crucial need to alter the behaviours of WP Labuan adolescents in order to prepare them for healthier adulthood. Poor dietary behaviours developed during adolescence may lead to diet related diseases in later years. Behaviour modification is the key recommendation suggested for improving healthy eating and lifestyle. It is necessary to improve dietary behaviour by encouraging them to consume nutritious foods such as fruits, vegetables, milk and milk products. This may be achieved through strengthening school-based nutrition interventions, using social marketing approach and mobilising families and communities into support. It

is crucial to extend school-based nutrition intervention programmes, such as *Program Hidangan Berkhasiat di Sekolah* (HiTS) to all schools. Aggressive promotion of healthy foods and the benefits of eating it should be made in all platforms, including social media. Intersectoral collaboration through various sectors is essential for the implementation of these strategies, so that nutrition programmes could be incorporated into their policies and improve access and availability of healthy foods in schools, food premises and the community.

**Table: Dietary Behaviour Trend among Adolescents in WP Labuan, Malaysia**

	NHMS 2012	NHMS 2017	NHMS 2022
Most of the time or always went hungry	Not reported	4.3	4.1
Fruits intake of at least twice daily	Not reported	50.7	35.5
Vegetables intake of at least three times daily	Not reported	42.3	25.6
Fruits and vegetables intake of at least five times daily	Not reported	28.7	15.9
Carbonated soft drinks consumption at least once daily	Not reported	52.4	46.5
Milk/milk products intake of at least two times daily	Not reported	35.2	25.7
Fast food intake of at least three days in the past seven days	Not reported	16.9	15.1

**3.3.7 References**

1. Fletcher EA, McNaughton SA, Crawford D, Cleland V, Della Gatta J, Hatt J, Dollman J, Timperio A. Associations between sedentary behaviours and dietary intakes among adolescents. *Public Health Nutr.* 2018 Apr;21(6):1115-1122
2. Mittal M, Jain V. Management of Obesity and Its Complications in Children and Adolescents. *Indian J Pediatr.* 2021 Dec;88(12):1222-1234
3. Kansra AR, Lakkunarajah S, Jay MS. Childhood and Adolescent Obesity: A Review. *Front Pediatr.* 2021 Jan 12;8:581461

**Table 3.3.1: Prevalence of adolescents in WP Labuan who most of the time or always went hungry in the past 30 days because there was not enough food in his/her home, 2022**

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
<b>WP LABUAN</b>	80	267	4.1	2.97	5.66
<b>Sex</b>					
Male	27	92	2.9	1.67	5.02
Female	53	175	5.2	4.00	6.85
<b>Form</b>					
Form 1	21	75	5.2	3.76	7.04
Form 2	24	83	5.6	3.09	9.77
Form 3	13	-	-	-	-
Form 4	10	-	-	-	-
Form 5	12	-	-	-	-
<b>Ethnicity</b>					
Malay	41	136	4.7	2.87	7.48
Chinese	8	-	-	-	-
Indian	0	-	-	-	-
Bumiputera Sabah	21	73	4.0	2.61	6.12
Bumiputera Sarawak	0	-	-	-	-
Others	10	-	-	-	-

- Prevalence with high RSE, not reported

**Table 3.3.2: Prevalence of fruit intake of at least twice daily in the past 30 days among adolescents in WP Labuan, 2022**

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
<b>WP LABUAN</b>	725	2306	35.5	34.12	36.89
<b>Sex</b>					
Male	367	1167	36.9	33.25	40.62
Female	358	1139	34.2	32.22	36.23
<b>Form</b>					
Form 1	144	506	34.6	30.10	39.30
Form 2	166	575	38.6	34.40	42.99
Form 3	122	427	32.2	26.75	38.23
Form 4	145	398	34.0	26.72	42.10
Form 5	148	401	38.2	35.52	40.90
<b>Ethnicity</b>					
Malay	325	1019	35.0	31.16	39.07
Chinese	78	261	33.5	25.15	43.12
Indian	2	-	-	-	-
Bumiputera Sabah	202	639	35.2	30.94	39.73
Bumiputera Sarawak	26	85	34.4	24.34	46.11
Others	92	294	40.5	33.03	48.36

- Prevalence with high RSE, not reported

**Table 3.3.3: Prevalence of vegetables intake of at least three times daily in the past 30 days among adolescents in WP Labuan, 2022**

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
<b>WP LABUAN</b>	519	1665	25.6	23.76	27.59
<b>Sex</b>					
Male	279	890	28.1	26.11	30.21
Female	240	775	23.3	21.02	25.68
<b>Form</b>					
Form 1	115	408	27.9	19.79	37.73
Form 2	98	336	22.6	17.85	28.11
Form 3	105	374	28.2	21.71	35.78
Form 4	102	282	24.1	18.19	31.22
Form 5	99	265	25.3	21.40	29.57
<b>Ethnicity</b>					
Malay	216	686	23.6	20.38	27.10
Chinese	61	207	26.6	24.22	29.17
Indian	1	-	-	-	-
Bumiputera Sabah	158	496	27.3	22.39	32.91
Bumiputera Sarawak	27	92	37.3	25.93	50.20
Others	56	179	24.7	18.47	32.17

- Prevalence with high RSE, not reported

**Table 3.3.4: Prevalence of fruits and vegetables intake of at least five times daily in the past 30 days among adolescent in WP Labuan, 2022**

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
<b>WP LABUAN</b>	322	1031	15.9	14.19	17.72
<b>Sex</b>					
Male	168	541	17.1	14.17	20.46
Female	154	490	14.7	13.62	15.89
<b>Form</b>					
Form 1	65	231	15.8	11.18	21.87
Form 2	66	223	15.0	11.31	19.62
Form 3	59	211	16.0	11.48	21.78
Form 4	62	173	14.8	9.30	22.66
Form 5	70	192	18.3	15.30	21.76
<b>Ethnicity</b>					
Malay	134	428	14.7	11.65	18.40
Chinese	38	129	16.6	13.07	20.76
Indian	1	-	-	-	-
Bumiputera Sabah	97	301	16.6	13.36	20.43
Bumiputera Sarawak	14	46	18.7	11.39	29.26
Others	38	123	16.9	10.69	25.72

- Prevalence with high RSE, not reported

Table 3.3.5: Prevalence of adolescents who did not consume fruit, vegetable or both in the past 30 days in WP Labuan, 2022

Socio-demographic characteristics	Never consume fruit				Never consume vegetable				Never consume fruit and vegetable			
	Unweighted count	Estimated population	Prevalence (%)	95 % CI Lower Upper	Unweighted count	Estimated population	Prevalence (%)	95 % CI Lower Upper	Unweighted count	Estimated population	Prevalence (%)	95 % CI Lower Upper
<b>WP LABUAN</b>	267	873	13.4	11.16 16.09	235	761	11.7	9.11 14.93	67	214	3.3	2.50 4.31
<b>Sex</b>												
Male	121	399	12.6	9.81 16.09	118	382	12.1	8.43 16.98	38	121	3.8	2.47 5.87
Female	146	474	14.2	11.36 17.65	117	379	11.4	8.78 14.62	29	93	2.8	2.09 3.69
<b>Form</b>												
Form 1	67	230	15.8	12.09 20.27	67	238	16.3	12.07 21.57	14	-	-	-
Form 2	55	193	13.0	10.04 16.62	54	188	12.6	8.78 17.88	15	54	3.6	2.42 5.41
Form 3	58	202	15.3	10.20 22.29	37	-	-	-	12	-	-	-
Form 4	48	140	12.0	8.77 16.10	42	115	9.9	7.57 12.73	12	33	2.8	1.58 4.98
Form 5	39	107	10.2	6.99 14.60	35	95	9.0	5.64 14.12	14	38	3.6	2.34 5.61
<b>Ethnicity</b>												
Malay	135	445	15.3	12.41 18.68	145	459	15.8	14.15 17.50	46	147	5.1	3.92 6.52
Chinese	18	62	8.0	5.43 11.57	13	45	5.8	3.34 9.79	2	-	-	-
Indian	1	-	-	-	0	-	-	-	0	-	-	-
Bumiputera Sabah	70	231	12.7	9.04 17.57	45	155	8.5	4.77 14.73	10	-	-	-
Bumiputera Sarawak	13	43	17.3	10.38 27.31	8	-	-	-	4	-	-	-
Others	30	89	12.3	7.65 19.25	24	75	10.3	7.29 14.44	5	-	-	-

- Prevalence with high RSE, not reported

**Table 3.3.6: Prevalence of carbonated soft drinks intake of at least once a day in the past 30 days among adolescents in WP Labuan, 2022**

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
<b>WP LABUAN</b>	943	3020	46.5	42.12	50.89
<b>Sex</b>					
Male	462	1475	46.6	40.27	53.03
Female	481	1545	46.4	41.23	51.59
<b>Form</b>					
Form 1	193	677	46.3	38.22	54.51
Form 2	217	756	50.8	47.10	54.46
Form 3	175	608	46.0	38.77	53.32
Form 4	201	560	47.8	36.37	59.46
Form 5	157	418	39.8	27.70	53.38
<b>Ethnicity</b>					
Malay	440	1388	47.7	44.27	51.09
Chinese	73	249	32.0	21.11	45.29
Indian	2	-	-	-	-
Bumiputera Sabah	269	868	47.8	39.46	56.24
Bumiputera Sarawak	43	138	55.7	43.50	67.19
Others	116	371	51.0	45.18	56.85

- Prevalence with high RSE, not reported

**Table 3.3.7: Prevalence of plain water intake of less than 6 glasses per day in the past 30 days among adolescents in WP Labuan, 2022**

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
<b>WP LABUAN</b>	916	2916	44.9	42.76	47.04
<b>Sex</b>					
Male	383	1203	38.0	34.83	41.31
Female	533	1713	51.4	48.18	54.65
<b>Form</b>					
Form 1	198	687	47.0	42.83	51.16
Form 2	205	701	47.1	40.21	54.01
Form 3	170	581	43.9	36.00	52.06
Form 4	184	515	44.0	37.51	50.66
Form 5	159	433	41.2	35.76	46.90
<b>Ethnicity</b>					
Malay	436	1355	46.6	42.94	50.21
Chinese	99	336	43.1	39.65	46.71
Indian	1	-	-	-	-
Bumiputera Sabah	244	789	43.5	39.08	47.97
Bumiputera Sarawak	39	125	50.6	42.02	59.11
Others	97	307	42.3	33.35	51.78

- Prevalence with high RSE, not reported

**Table 3.3.8: Prevalence of milk and milk products intake of at least two servings per day in the past 30 days among adolescents in WP Labuan, 2022**

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
<b>WP LABUAN</b>	514	1668	25.7	22.97	28.56
<b>Sex</b>					
Male	241	782	24.7	21.07	28.75
Female	273	885	26.6	21.83	31.93
<b>Form</b>					
Form 1	106	376	25.7	20.45	31.70
Form 2	127	438	29.4	22.93	36.87
Form 3	100	356	26.9	19.59	35.64
Form 4	98	276	23.6	18.35	29.82
Form 5	83	222	21.1	18.87	23.56
<b>Ethnicity</b>					
Malay	222	710	24.4	19.81	29.66
Chinese	67	226	29.1	20.37	39.68
Indian	0	-	-	-	-
Bumiputera Sabah	135	439	24.2	19.03	30.15
Bumiputera Sarawak	25	80	32.2	25.54	39.67
Others	65	212	29.3	21.65	38.25

- Prevalence with high RSE, not reported

**Table 3.3.9: Prevalence of fast-food intake of at least three days in the past 7 days among adolescents in WP Labuan, 2022**

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
<b>WP LABUAN</b>	304	982	15.1	13.30	17.12
<b>Sex</b>					
Male	119	385	12.2	9.69	15.17
Female	185	597	17.9	15.68	20.38
<b>Form</b>					
Form 1	75	264	18.1	14.21	22.69
Form 2	58	198	13.3	9.55	18.16
Form 3	59	207	15.6	10.54	22.48
Form 4	58	166	14.2	10.77	18.50
Form 5	54	147	14.0	9.96	19.35
<b>Ethnicity</b>					
Malay	140	445	15.3	11.26	20.45
Chinese	25	85	10.9	7.35	15.98
Indian	0	-	-	-	-
Bumiputera Sabah	86	282	15.5	12.40	19.32
Bumiputera Sarawak	15	48	19.5	10.14	34.27
Others	38	121	16.6	12.02	22.51

- Prevalence with high RSE, not reported

### 3.4 Nutritional Status

Contributors: Ahmad Ali Zainuddin, Lalitha Palaniveloo, Khairul Hasnan Amali, Siti Adibah Ab. Halim, Ainan Nasrina Ismail

#### 3.4.1 Introduction

Adolescence is a unique phase of human development for individuals between the ages of 10 and 19 years old, as it caters to rapid growth, as well as sexual and behavioural changes. Good nutrition during adolescence is critical to address current nutritional needs and to fill nutrient gaps that have occurred during childhood<sup>1</sup>. The nutritional status of adolescents is assessed using anthropometric measurements (weight and height) and interpreted using WHO 2007 Growth Reference Data for 5-19 years<sup>2</sup>. The indicators include stunting, thinness, overweight and obesity.

#### 3.4.2 Objectives

- i. To determine the prevalence of stunting among adolescents
- ii. To determine the prevalence of thinness among adolescents
- iii. To determine the prevalence of overweight and obesity among adolescents

#### 3.4.3 Variable definitions

- **Body mass index (BMI):** commonly used to determine weight status. BMI is calculated by dividing a person's weight in kilograms by the square of height in meters.
- **Height for age z-score (HAZ):** an index used to assess how a child's height compares to the expected height of a healthy child of the same age and sex based on the WHO 2007 Growth reference data for 5-19 years.
- **BMI for age z-score (BAZ):** an index used to assess BMI is age- and sex-specific compares to the BMI of a healthy child of the same age and sex based on the WHO 2007 Growth reference data for 5-19 years.
- **Stunting:** Those who have their HAZ more than two standard deviations below the WHO Child Growth Standards median (<-2SD).
- **Thinness:** Those who have their BAZ more than two standard deviations below the WHO Child Growth Standards median (<-2SD).
- **Overweight:** Those who have their BAZ is more than one standard deviations to two standard deviations above the WHO Child Growth Standards median (>+1SD to ≤+2SD).
- **Obesity:** Those who have their BAZ more than two standard deviations above the WHO Child Growth Standards median (>+2SD).

#### 3.4.4 Findings

##### Height-for-Age z-score

The prevalence of stunting among adolescents was 7.8% (95% CI: 5.86, 10.38). Females [9.2%, (95% CI: 6.79, 12.24)] showed higher prevalence compared to males [6.4%, (95% CI: 3.99, 10.24)]. (Table 3.4.1).

##### BMI -for-Age z-score

According to the WHO 2007 Growth Reference Data for 5-19 years, the prevalence of thinness among adolescents was 6.1% (95% CI: 4.37, 8.33). The data showed that the prevalence of thinness was significantly higher among males [7.5%, (95% CI: 4.99, 11.22)] compared to females [4.6%, (95% CI: 3.11, 6.88)]. (Table 3.4.2).

##### Overweight and Obesity

The prevalence of overweight was 17.11% (95% CI: 15.29, 19.10). Comparing the sexes, females had a higher prevalence at 17.4% (95% CI: 14.92, 20.11)] compared to males [16.9% (95% CI: 13.30, 21.12)]. (Table 3.4.3). For obesity, the prevalence was 15.9% (95% CI: 13.25, 18.93). Between the sexes, males had a higher prevalence of obesity at 19.9% (95% CI: 15.30, 25.47) compared to females [12.0% (95% CI: 10.53, 13.73)]. (Table 3.4.3).

#### 3.4.5 Discussion / Conclusion

Overall, the prevalence of overweight and obesity totalling 33% was higher than thinness (6.1%) and stunting (7.8%). It can be concluded that adolescents in WP Labuan is facing a dual burden of malnutrition, with overnutrition appearing to be a larger problem than undernutrition.

#### 3.4.6 Recommendations

Based on the findings, the integration of targeted interventions and policies is required to simultaneously address both undernutrition and the increasing rates of overweight and obesity among adolescents. Evidence-based nutrition-sensitive interventions, inclusive of diet counselling and nutrition education provided through school-based platforms, adolescent youth centres/ peer education and technology-based platforms should be strengthened. A comprehensive intervention such as MyBFF@school (an intervention consisting of nutrition education, physical activity and motivational component) could be implemented nationwide with the support of the Ministry of Education.

#### 3.4.7 References

1. Das JK, Salam RA, Thornburg KL, et al. Nutrition in adolescents: physiology, metabolism, and nutritional needs. *Ann. N. Y. Acad. Sci.* 2017 Apr;1393(1):21-33
2. World Health Organization 2007. Growth reference data for 5-19 years. Geneva: WHO

Table 3.4.1: Prevalence of stunting (HAZ &lt;-2SD) among adolescents in WP Labuan, 2022

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
<b>WP LABUAN</b>	167	506	7.8	5.86	10.38
<b>Sex</b>					
Male	70	204	6.4	3.99	10.24
Female	97	302	9.2	6.79	12.24
<b>Form</b>					
Form 1	18	-	-	-	-
Form 2	24	79	5.3	3.65	7.73
Form 3	29	96	7.3	3.48	14.71
Form 4	57	162	13.9	10.84	17.63
Form 5	39	105	10.0	7.46	13.28
<b>Ethnicity</b>					
Malay	91	267	9.2	6.56	12.80
Chinese	9	-	-	-	-
Indian	0	-	-	-	-
Bumiputera Sabah	45	139	7.7	5.02	11.67
Bumiputera Sarawak	3	-	-	-	-
Others	19	-	-	-	-

- Prevalence with high RSE, not reported

Table 3.4.2: Prevalence of thinness (BAZ &lt;-2SD) among adolescents in WP Labuan, 2022

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
<b>WP LABUAN</b>	126	391	6.1	4.37	8.33
<b>Sex</b>					
Male	79	238	7.5	4.99	11.22
Female	47	153	4.6	3.11	6.88
<b>Form</b>					
Form 1	23	80	5.6	3.38	9.03
Form 2	23	83	5.6	3.75	8.30
Form 3	13	46	3.5	2.00	5.99
Form 4	38	106	9.0	5.48	14.58
Form 5	29	76	7.3	4.39	11.89
<b>Ethnicity</b>					
Malay	66	204	7.0	5.62	8.77
Chinese	7	-	-	-	-
Indian	0	-	-	-	-
Bumiputera Sabah	29	-	-	-	-
Bumiputera Sarawak	5	-	-	-	-
Others	19	59	8.1	4.72	13.55

- Prevalence with high RSE, not reported



Table 3.4.3: Prevalence of overweight (BAZ >+1SD to ≤+2SD) and obesity (BAZ >+2SD) among adolescents in WP Labuan, 2022

Socio-demographic characteristics	Overweight (>+1SD to ≤+2SD)					Obese (>+2SD)				
	Unweighted count	Estimated population	Prevalence (%)	95 % CI Lower	Upper	Unweighted count	Estimated population	Prevalence (%)	95 % CI Lower	Upper
<b>WP LABUAN</b>	344	1104	17.1	15.29	19.10	313	1025	15.9	13.25	18.93
<b>Sex</b>										
Male	167	532	16.9	13.30	21.12	190	628	19.9	15.30	25.47
Female	177	573	17.4	14.92	20.11	123	397	12.0	10.53	13.73
<b>Form</b>										
Form 1	60	213	14.7	11.20	19.15	85	303	21.0	14.37	29.62
Form 2	76	-	-	-	-	72	258	17.5	13.02	23.12
Form 3	67	237	17.9	12.36	25.33	54	187	14.2	9.54	20.52
Form 4	75	212	18.2	12.34	26.02	54	149	12.8	8.65	18.49
Form 5	66	177	16.9	10.63	25.78	48	127	12.2	8.47	17.18
<b>Ethnicity</b>										
Malay	156	499	17.2	13.60	21.59	149	493	17.0	13.37	21.36
Chinese	47	156	20.2	14.49	27.35	26	86	11.1	7.36	16.52
Indian	1	-	-	-	-	1	-	-	-	-
Bumiputera Sabah	96	306	17.1	13.19	21.77	85	278	15.5	10.63	22.03
Bumiputera Sarawak	10	-	-	-	-	11	38	15.4	9.13	24.81
Others	19	59	8.1	4.72	13.55	41	-	-	-	-

- Prevalence with high RSE, not reported

## 3.5 Drug Use

Contributors: Thamil Arasu Saminathan, Hamizatul Akmal Abd Hamid, Muhammad Fadhli Mohd Yusoff, Tania Gayle Robert, Maznieda Mahjom, Hasimah Ismail, Mohd Haniff Bin Bistari, Halizah Mat Rifin, Mohamad Salleh Abdul Ghani, Norli Abdul Jabbar, Rushidi Abd Rashid

### 3.5.1 Introduction

According to the World Drug Report 2022 (WDR 2022) by the United Nations Office on Drugs and Crime (UNODC), an estimated 284 million people had used drugs within the previous year, which accounts for a 26% increase over the previous decade.<sup>1</sup> Drug use accounts for 5% of all substance related death and 9% of substance-use-related DALYs. Despite the report showing that young people continue to use more drugs than adults, it was found that drug use by adolescents decreased during the COVID-19 pandemic, which coincided with the lockdown periods.<sup>1</sup> Marijuana or cannabis remains the world's most widely used drug, with an annual prevalence of 4% of the adult population, or an estimated 209 million users in the past year.<sup>1</sup> Amphetamines remain the second most commonly used drug worldwide, with an estimated 34 million in 2020, representing 0.7% of the global population. NHMS 2019 showed that marijuana is the highest taken in Malaysia, followed by kratom.<sup>2</sup> Based on the statistics provided by the National Anti-Drug Agency in 2020 showed that ATS is the most commonly used drug among adolescents in Malaysia aged 13 to 18 years old, followed by marijuana and opiate.<sup>3</sup> In this survey, we have added new questions, which include kratom and inhalant, to get baseline data on adolescent usage in Malaysia, as we don't have preliminary national data on these drugs.

### 3.5.2 Objectives

- i. To determine the prevalence and sociodemographic characteristics of ever and current drug use among adolescents
- ii. To determine the prevalence and sociodemographic characteristics of ever and current marijuana use among adolescents in Malaysia
- iii. To determine the prevalence and sociodemographic characteristics of ever and current amphetamines or methamphetamines use among adolescents in Malaysia
- iv. To determine the prevalence and sociodemographic characteristics of ever and current inhalant use among adolescents in Malaysia
- v. To determine the prevalence and sociodemographic characteristics of ever and current kratom use among adolescents in Malaysia
- vi. To identify the age of initiation and the sources of obtaining drugs among adolescents in Malaysia

### 3.5.3 Variable Definitions

- **Drug use:**
  - i. **2017** definition: taking heroin, morphine, glue, amphetamine, or methamphetamines (ecstasy, syabu, ice), marijuana (except prescribed medicine).
  - ii. **2022** definition: taking opiates, amphetamine-type stimulants, marijuana, psychotropic pill, cocaine, inhalant and others (depressants, hallucinogens).
- **Ever drug use:** adolescents who had a history of drug use in their lifetime
- **Current drug use:** adolescents who used drugs in the past 30 days
- **Ever marijuana use:** adolescents who had a history of marijuana use in their lifetime
- **Current marijuana use:** adolescents who used marijuana in the past 30 days
- **Ever amphetamine or methamphetamine use:** adolescents with a history of amphetamine or methamphetamines use in their lifetime
- **Ever inhalant use:** adolescents who had a history of inhalant use in their lifetime
- **Current inhalant use:** adolescents who used an inhalant in the past 30 days
- **Ever kratom use:** adolescents who had a history of kratom use in their lifetime
- **Current kratom use:** adolescents who used kratom in the past 30 days

### 3.5.4 Findings

Overall, 3.9% (95% CI: 2.96, 5.11) of adolescents reported that they had ever used drug during their lifetime and it was significantly higher among males [5.0% (95%CI: 3.64, 6.78)] as compared to females [2.9% (95%CI: 1.78, 4.56)] (**Table 3.5.1**). The prevalence of current drug users was 2.1% (95%CI: 1.25, 3.40), males [3.1% (95%CI: 1.73, 5.59)] (**Table 3.5.2**). Overall, 2.4% (95% CI: 1.66, 3.56) of adolescents reported had ever used inhalant during their lifetime. Males was also significantly higher [2.9% (95% CI: 1.61, 5.17)] than females [2.0% (95% CI: 1.23, 3.20)]. Overall, 0.6% (95% CI: 0.31, 0.17) of female adolescents reported had current used inhalant in the past 30 days. (**Table 3.5.3**). Among ever drug users, 79.2% (95% CI: 39.08, 95.78) of them had initiated before the age of 14 years old (**Table 3.5.4**).

### 3.5.5 Discussion / Conclusion

Unlike other drugs, this survey found that inhalant use was most common among male adolescents, which tended to decline as they grew older. The early use of inhalants in both sexes reflects inhalants as the "kids' drug", which are available legally.

### 3.5.6 Recommendations

The effectiveness of drug education at primary schools plays a vital role in reducing current drug use prevalence in 2022 compared to 2017. Drug prevention among adolescents should be improved and regularly reviewed to meet the ever-changing trend of drug use locally and globally. New strategies and approaches can be developed to address issues of inhalant and kratom use among adolescents by focusing more on the danger of inhalant and kratom use. An adolescent who has been exposed to and involved in drug abuse must be given access to treatment and provided educational opportunities, vocational skills training and other socioeconomic support needed. School drug prevention programs developed for adolescents can be integrated with technology, such as web-based intervention, to make it more interesting in increasing awareness and help build self-resilience among adolescents through:

- Conducting early detection of an adolescent with problems or who are at risk of drug abuse
- Increasing the knowledge, understanding and awareness of the dangers of drug abuse
- Increasing life skills such as assertiveness, coping skills and stress management
- Reinforcing positive attitudes and healthy lifestyles among school children

### 3.5.7 References

1. United Nations Office on Drugs and Crime (UNODC), World Drug Report 2022; Booklet 1; Executive Summary and Policy Implication (ISBN: 9789211483758) <https://www.unodc.org/unodc/en/data-and-analysis/world-drug-report-2022.html>
2. Institute for Public Health (IPH) 2020. The National Health and Morbidity Survey 2019: NCD. Kuala Lumpur: Ministry of Health Malaysia
3. Bahagian Dasar, Perancangan dan Penyelidikan, Agensi Antidadah Kebangsaan, Kementerian Dalam Negeri; Info Dadah Siri 1/2020; Penyalahgunaan Dadah Dalam Kalangan Remaja; 4 JUN 2020

Table 3.5.1: Prevalence of ever drug use among adolescents in WP Labuan, 2022

Socio-demographic characteristics	Ever used drug* based on 2017				Ever used drug**			
	Unweighted count	Estimated population	Prevalence (%)	95 % CI Lower Upper	Unweighted count	Estimated population	Prevalence (%)	95 % CI Lower Upper
<b>WP LABUAN</b>	64	210	3.2	2.28 4.56	79	253	3.9	2.96 5.11
<b>Sex</b>								
Male	39	127	4.0	2.62 6.16	50	158	5.0	3.64 6.78
Female	25	82	2.5	1.55 3.93	29	95	2.9	1.78 4.56
<b>Form</b>								
Form 1	18	-	-	- -	20	-	-	- -
Form 2	12	-	-	- -	14	-	-	- -
Form 3	15	52	3.9	2.25 6.79	16	56	4.2	2.49 7.07
Form 4	9	-	-	- -	14	-	-	- -
Form 5	10	27	2.6	1.52 4.48	15	40	3.8	3.06 4.79
<b>Ethnicity</b>								
Malay	24	75	2.6	1.59 4.16	33	100	3.4	2.32 5.09
Chinese	8	-	-	- -	10	36	4.7	2.91 7.45
Indian	0	-	-	- -	0	-	-	- -
Bumiputera Sabah	16	53	2.9	1.87 4.47	17	55	3.0	2.05 4.48
Bumiputera Sarawak	2	-	-	- -	4	-	-	- -
Others	14	47	6.4	3.32 12.05	15	50	6.8	3.49 12.91

- Prevalence with high RSE, not reported

\*Drug includes heroin, morphine, glue, amphetamine, ecstasy, methamphetamine, ice and marijuana.

\*\*Drug includes opiate, amphetamine, marijuana, psychotropic pill, cocaine, inhalant, kratom and others.

Table 3.5.2: Prevalence of current drug use among adolescents in WP Labuan, 2022

Socio-demographic characteristics	Current used drug* based on 2017				Current used drug** based on 2022			
	Unweighted count	Estimated population	Prevalence (%)	95 % CI Lower Upper	Unweighted count	Estimated population	Prevalence (%)	95 % CI Lower Upper
<b>WP LABUAN</b>	35	113	1.7	1.06 2.84	42	135	2.1	1.25 3.40
<b>Sex</b>								
Male	26	84	2.6	1.55 4.47	31	99	3.1	1.73 5.59
Female	9	-	-	- -	11	-	-	- -
<b>Form</b>								
Form 1	15	-	-	- -	16	-	-	- -
Form 2	4	-	-	- -	6	-	-	- -
Form 3	5	-	-	- -	6	-	-	- -
Form 4	7	-	-	- -	7	-	-	- -
Form 5	4	-	-	- -	7	-	-	- -
<b>Ethnicity</b>								
Malay	16	-	-	- -	20	-	-	- -
Chinese	5	-	-	- -	6	23	2.9	1.52 5.56
Indian	0	-	-	- -	0	-	-	- -
Bumiputera Sabah	7	-	-	- -	8	-	-	- -
Bumiputera Sarawak	3	-	-	- -	3	-	-	- -
Others	4	-	-	- -	5	-	-	- -

- Prevalence with high RSE, not reported

\*Drug includes heroin, morphine, glue, amphetamine, ecstasy, methamphetamine, ice and marijuana.

\*\*Drug includes opiate, amphetamine, marijuana, psychotropic pill, cocaine, inhalant, kratom and others.

Table 3.5.3: Prevalence of inhalant use among adolescents in WP Labuan, 2022

Socio-demographic characteristics	Ever used inhalant in a lifetime				Current used inhalant in the past 30 days			
	Unweighted count	Estimated population	Prevalence (%)	95 % CI Lower Upper	Unweighted count	Estimated population	Prevalence (%)	95 % CI Lower Upper
<b>WP LABUAN</b>	48	158	2.4	1.66 3.56	21	-	-	- -
<b>Sex</b>								
Male	28	92	2.9	1.61 5.17	15	-	-	- -
Female	20	66	2.0	1.23 3.20	6	20	0.6	0.31 1.17
<b>Form</b>								
Form 1	14	-	-	- -	10	-	-	- -
Form 2	12	-	-	- -	4	-	-	- -
Form 3	9	-	-	- -	1	-	-	- -
Form 4	7	-	-	- -	5	-	-	- -
Form 5	6	-	-	- -	1	-	-	- -
<b>Ethnicity</b>								
Malay	17	52	1.8	0.97 3.31	7	-	-	- -
Chinese	7	-	-	- -	4	-	-	- -
Indian	0	-	-	- -	0	-	-	- -
Bumiputera Sabah	11	-	-	- -	5	-	-	- -
Bumiputera Sarawak	2	-	-	- -	2	-	-	- -
Others	11	-	-	- -	3	-	-	- -

- Prevalence with high RSE, not reported

Table 3.5.6: Prevalence of first use of drug before the age of 14 years among ever used drug among adolescents in WP Labuan, 2022

Prevalence	Unweighted count	Estimated population	Prevalence (%)	95 % CI Lower Upper
Yes	8	28	79.2	39.08 95.78
No	3	-	-	- -

- Prevalence with high RSE, not reported

## 3.6 Oral and Hand Hygiene

Contributors: Fazila Haryati Ahmad, Rafidah Ali, Chan Yee Mang, Mohd Hatta Abdul Mutalip, Nik Adilah Shahein, Norzawati Yoep, Nurulasmak Mohamed, Saidatul Norbaya Buang, Nik Daliana Nik Farid, Annapurny Venkiteswaran

### 3.6.1 Introduction

Oral health is integral to general health as it promotes a positive quality of life and social self-confidence. Currently, oral diseases affect close to 3.5 billion people worldwide, and their prevalence is noted to be increasing globally. A resolution on oral health in 2021 by the WHO recommends a more preventive approach towards oral health, including oral health promotions at schools<sup>1</sup>. Empowering good oral hygiene habits during adolescence is important in sustaining this behaviour into adulthood<sup>2</sup>. Therefore, early and adequate plaque control is key in preventing oral health diseases such as dental caries and periodontal diseases, which may affect school performance and attendance, as well as permanent dental problems in adulthood. This can be achieved via regular tooth brushing with fluoridated toothpaste, dental flossing, tongue cleaning, and a minimum yearly dental check-up<sup>3</sup>. Appropriate hand hygiene practices using soap, especially before eating and after using the toilet, are protective against a multitude of infections. These practices will enable adolescents to thrive and contribute actively to learning and reduce the rate of absenteeism<sup>4</sup>. Assessing practices on good hand washing among adolescents will help detect at-risk groups among school attendees<sup>4</sup>.

### 3.6.2 Objectives

#### 3.6.2.1 General objective

To determine the prevalence of oral and hand hygiene behaviour among adolescents in Malaysia.

#### 3.6.2.2 Specific objectives for oral hygiene

To describe the prevalence of:

- i. Self-oral health perception
- ii. Tooth brushing frequency in the past 30 days
- iii. Tongue cleaning practice
- iv. Fluoridated toothpaste usage
- v. Dental floss usage
- vi. Timing of the last visit to a dentist or dental nurse
- vii. Having missed class or not participating in online learning due to toothache in the past 12 months
- viii. Avoidance of smile or laughing due to the appearance of their teeth

#### 3.6.2.3 Specific objectives for hand washing

To describe the prevalence of:

- i. Hand washing with soap in the past 30 days
- ii. Hand washing before eating in the past 30 days
- iii. Hand washing after using the toilet in the past 30 days
- iv. Hand washing method before eating at school in the past 30 days

### 3.6.3 Variable definitions

- **Clean or brush teeth:** Regular tooth brushing using toothbrush and toothpaste to keep the mouth, teeth and gums clean and healthy
- **Last saw a dentist or dental nurse:** Seen a dentist or dental nurse for a check-up, scaling or other dental treatment

### 3.6.4 Findings

2.7% (95%CI: 2.05, 3.51) of adolescents in WP Labuan perceived their oral health as poor or very poor (**Table 3.6.1**). 85.7% (95%CI: 83.33, 87.82) brushed their teeth twice daily which was significantly higher in females [90.8% (95%CI: 89.03, 92.24)] (**Table 3.6.2**), 4.6% (95% CI: 3.45, 6.20) never performed daily tongue cleaning (**Table 3.6.3**), 43.3% (95%CI: 40.08, 46.55) reported not knowing whether their toothpaste contained fluoride (**Table 3.6.4**) and only 24.1% (95%CI: 22.31, 26.05) used dental floss for cleaning their teeth (**Table 3.6.5**). Only 32.8% (95%CI: 25.49, 40.98) reported to have their last dental visit in the past 12 months (**Table 3.6.6**), 9.0% (95% CI: 6.37, 12.58) had toothache in the past 12 months and had missed class or not participated with online learning (PdPR) (**Table 3.6.7**), 33.1% (95%CI: 28.35, 38.18) reported that they had avoided smiling or laughing due to the appearance of their teeth which was significantly higher among females [41.7% (95%CI: 35.74, 47.99)] (**Table 3.6.8**). The prevalence of WP Labuan adolescents who used soap most of the time or always was 72.8% (95%CI: 69.04, 76.27) (**Table 3.6.9**). About 84.9% (95%CI: 82.51, 87.05) and 87.5% (95%CI: 85.39, 89.35) of adolescents washed their hands most of the time or always before eating (**Table 3.6.10**) or after using toilet respectively (**Table 3.6.11**). Only 48.2% (95%CI: 42.17, 54.37) washed hands with running water before eating at school (**Table 3.6.12**).

### 3.6.5 Discussion / Conclusion

In general, oral hygiene behaviour among adolescents may be related to the COVID-19 pandemic that reduces school dental programs and attendance for dental checkups, which in turn may cause inadequate knowledge regarding best oral health practice among adolescents. Ironically, hand hygiene practice appear to be good which may also reflect the effect of COVID-19 pandemic that promotes frequent handwashing practice in general.

### 3.6.6 Recommendations

Taking cognizance of these findings, there is a need for continuous emphasis on promoting good personal oral and hand hygiene among adolescents through knowledge, attitude and behavioural improvements with these following recommendations:

- Oral health education at schools need to deliver captivating methods that can be easily assimilated into the adolescents' daily school and home routines which will enhance their retention of oral health care knowledge. Effective oral health education should be regularly revised, updated and tailored specifically for young adults to improve and empower their decision making in maintaining good oral health. Adolescents at high risk of developing oral diseases should be identified early and oral health intervention delivered and tailored to these targeted groups to enhance engagement and personalisation of oral care needs.
- Interventions to promote hand washing need to be tailored to the adolescent's understanding and relevant social norms to trigger and reinforce good and ideal handwashing practice and habit formation according to their environment and social situations. Schools should have policies that inculcate good handwashing behavior including adequate infrastructures to support regular and ideal handwashing practices, and readily available information visual aids about the correct hand washing technique at key times and places in schools. School health education unit with regular hygiene education programs should be included in the formal and non-formal curricular, preferably starting from pre and primary schools. Further studies are needed to develop an understanding and knowledge gap of the cultural context of handwashing habits in various ethnic and cultural groups.

### 3.6.7 References

1. World Health Organization. (2022, October 10). Fact Sheets on Oral Health: WHO Response. Retrieved from <https://www.who.int/news-room/fact-sheets/detail/oral-health>
2. Calderon S, Mallory C. Look at My Pearly White Teeth: Exploring Adolescents' Oral Health Behavior. *Public Health Nurs.* 2018; 1-8
3. Malaysian Dental Association. General Oral Health Care, Question 3: Teeth for Life? [Online]. 2020 [Updated 18 April 2020]. <https://web.mda.org.my/questions-3-teeth-for-life-contributed-by-prof-dr-ishak-abdul-razak/>. Accessed on 13 October 2022
4. Habib R. Effect of Hand Washing Practices and Prevalence of Related Diseases among Primary School Children in Tehsil Lalian, District Chiniot, Pakistan. *PriMera Scientific Medicine and Public Health.* 2022;1:15-26



**Table 3.6.1: Prevalence of poor or very poor perception of oral health among adolescents in WP Labuan, 2022**

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
<b>WP LABUAN</b>	52	175	2.7	2.05	3.51
<b>Sex</b>					
Male	27	93	2.9	2.01	4.25
Female	25	82	2.5	1.35	4.44
<b>Form</b>					
Form 1	14	48	3.3	2.24	4.83
Form 2	17	60	4.1	2.97	5.53
Form 3	11	-	-	-	-
Form 4	4	-	-	-	-
Form 5	6	-	-	-	-
<b>Ethnicity</b>					
Malay	24	79	2.7	1.95	3.76
Chinese	6	-	-	-	-
Indian	0	-	-	-	-
Bumiputera Sabah	14	48	2.6	1.39	5.00
Bumiputera Sarawak	1	-	-	-	-
Others	7	-	-	-	-

- Prevalence with high RSE, not reported

**Table 3.6.2: Prevalence of teeth brushing 2 times a day in the past 30 days among adolescents in WP Labuan, 2022**

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
<b>WP LABUAN</b>	1745	5569	85.7	83.33	87.82
<b>Sex</b>					
Male	802	2545	80.4	77.30	83.22
Female	943	3024	90.8	89.03	92.24
<b>Form</b>					
Form 1	341	1196	81.7	76.11	86.28
Form 2	371	1276	85.7	78.18	90.93
Form 3	330	1150	86.9	81.77	90.68
Form 4	366	1031	88.0	83.11	91.67
Form 5	337	917	87.3	73.36	94.51
<b>Ethnicity</b>					
Malay	797	2513	86.3	82.52	89.42
Chinese	192	641	82.3	74.33	88.23
Indian	4	14	79.2	18.40	98.47
Bumiputera Sabah	476	1528	84.2	78.90	88.33
Bumiputera Sarawak	69	223	89.8	77.10	95.82
Others	207	650	89.5	85.77	92.41

Table 3.6.3: Prevalence of never did tongue cleaning among adolescents in WP Labuan, 2022

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
<b>WP LABUAN</b>	93	301	4.6	3.45	6.20
<b>Sex</b>					
Male	62	198	6.3	4.60	8.48
Female	31	103	3.1	1.71	5.51
<b>Form</b>					
Form 1	25	88	6.0	3.04	11.68
Form 2	19	67	4.5	2.54	7.91
Form 3	13	-	-	-	-
Form 4	23	64	5.4	3.11	9.36
Form 5	13	33	3.2	1.61	6.09
<b>Ethnicity</b>					
Malay	42	127	4.4	2.92	6.46
Chinese	21	73	9.4	6.27	13.93
Indian	1	-	-	-	-
Bumiputera Sabah	17	59	3.3	1.73	6.06
Bumiputera Sarawak	1	-	-	-	-
Others	11	35	4.8	2.87	8.08

- Prevalence with high RSE, not reported

Table 3.6.4: Prevalence of did not know if their toothpaste is fluoridated among adolescents in WP Labuan, 2022

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
<b>WP LABUAN</b>	868	2810	43.3	40.08	46.55
<b>Sex</b>					
Male	427	1383	43.8	40.16	47.49
Female	441	1426	42.8	37.53	48.25
<b>Form</b>					
Form 1	184	639	43.7	37.18	50.43
Form 2	214	738	49.8	42.61	56.91
Form 3	170	610	46.1	35.02	57.57
Form 4	180	504	43.1	34.26	52.30
Form 5	120	318	30.3	23.31	38.34
<b>Ethnicity</b>					
Malay	381	1221	42.0	36.93	47.24
Chinese	126	431	55.3	47.55	62.87
Indian	0	-	-	-	-
Bumiputera Sabah	238	765	42.2	37.39	47.20
Bumiputera Sarawak	34	109	43.9	27.23	62.10
Others	89	283	39.0	31.00	47.67

- Prevalence with high RSE, not reported

**Table 3.6.5: Prevalence of use of dental floss among adolescents in WP Labuan, 2022**

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
<b>WP LABUAN</b>	489	1568	24.1	22.31	26.05
<b>Sex</b>					
Male	211	673	21.2	18.96	23.74
Female	278	895	26.9	23.51	30.52
<b>Form</b>					
Form 1	108	381	26.0	19.35	34.03
Form 2	97	329	22.1	17.54	27.37
Form 3	86	302	22.8	15.77	31.87
Form 4	98	280	24.0	20.22	28.15
Form 5	100	276	26.2	18.53	35.78
<b>Ethnicity</b>					
Malay	212	666	22.9	20.72	25.22
Chinese	64	215	27.7	22.91	32.99
Indian	0	-	-	-	-
Bumiputera Sabah	144	468	25.8	21.25	30.85
Bumiputera Sarawak	21	69	27.6	17.52	40.66
Others	48	150	20.6	16.79	25.10

- Prevalence with high RSE, not reported

**Table 3.6.6: Prevalence of last dental visit in the past 12 months among adolescents in WP Labuan, 2022**

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
<b>WP LABUAN</b>	685	2129	32.8	25.49	40.98
<b>Sex</b>					
Male	334	1032	32.6	22.19	45.08
Female	351	1097	32.9	24.13	43.09
<b>Form</b>					
Form 1	119	416	28.4	22.35	35.39
Form 2	124	422	28.3	20.01	38.42
Form 3	126	430	32.4	20.55	47.14
Form 4	144	404	34.5	21.44	50.41
Form 5	172	458	43.6	29.34	59.02
<b>Ethnicity</b>					
Malay	311	937	32.2	22.55	43.62
Chinese	79	262	33.6	21.38	48.49
Indian	2	-	-	-	-
Bumiputera Sabah	196	619	34.1	24.92	44.69
Bumiputera Sarawak	33	104	41.8	26.71	58.59
Others	64	200	27.5	19.76	36.97

- Prevalence with high RSE, not reported

Table 3.6.7: Prevalence of having missed classes or online learning among adolescents with toothache in WP Labuan, 2022

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
<b>WP LABUAN</b>	94	308	9.0	6.37	12.58
<b>Sex</b>					
Male	48	162	9.4	5.88	14.75
Female	46	145	8.6	5.40	13.35
<b>Form</b>					
Form 1	25	88	10.8	5.41	20.23
Form 2	20	-	-	-	-
Form 3	14	48	7.0	3.41	13.83
Form 4	14	41	6.8	3.92	11.53
Form 5	21	57	10.3	6.82	15.40
<b>Ethnicity</b>					
Malay	42	138	9.1	5.70	14.10
Chinese	7	-	-	-	-
Indian	0	-	-	-	-
Bumiputera Sabah	23	-	-	-	-
Bumiputera Sarawak	2	-	-	-	-
Others	20	-	-	-	-

- Prevalence with high RSE, not reported

Table 3.6.8: Prevalence of avoidance of smiling due to teeth appearance among adolescents in WP Labuan, 2022

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
<b>WP LABUAN</b>	664	2149	33.1	28.35	38.18
<b>Sex</b>					
Male	235	759	24.0	20.17	28.24
Female	429	1391	41.7	35.74	47.99
<b>Form</b>					
Form 1	131	458	31.3	26.79	36.18
Form 2	157	531	35.7	27.25	45.05
Form 3	144	494	37.3	27.73	47.96
Form 4	127	373	31.8	24.72	39.88
Form 5	105	294	28.0	21.47	35.67
<b>Ethnicity</b>					
Malay	301	957	32.9	28.97	37.02
Chinese	78	264	33.9	22.59	47.46
Indian	2	-	-	-	-
Bumiputera Sabah	192	625	34.4	26.91	42.83
Bumiputera Sarawak	26	86	34.8	22.90	48.87
Others	65	210	28.9	20.28	39.42

- Prevalence with high RSE, not reported

**Table 3.6.9: Prevalence of using soap most of the time or always during handwashing in the past 30 days among adolescents in WP Labuan, 2022**

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95% CI	
				Lower	Upper
<b>WP LABUAN</b>	1485	4730	72.8	69.04	76.27
<b>Sex</b>					
Male	681	2149	67.9	61.27	73.90
Female	804	2581	77.5	72.02	82.10
<b>Form</b>					
Form 1	282	993	67.9	59.87	74.96
Form 2	302	1045	70.2	60.25	78.54
Form 3	267	931	70.3	61.68	77.72
Form 4	312	885	75.6	65.23	83.64
Form 5	322	876	83.4	73.03	90.32
<b>Ethnicity</b>					
Malay	662	2073	71.2	66.11	75.81
Chinese	174	577	74.1	64.60	81.80
Indian	4	14	79.2	18.40	98.47
Bumiputera Sabah	425	1362	75.0	67.60	81.17
Bumiputera Sarawak	58	192	77.5	63.62	87.10
Others	162	513	70.6	59.38	79.80

**Table 3.6.10: Prevalence of handwashing most of the time or always before eating in the past 30 days among adolescents in WP Labuan, 2022**

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
<b>WP LABUAN</b>	1732	5517	84.9	82.51	87.05
<b>Sex</b>					
Male	840	2660	84.0	80.05	87.38
Female	892	2857	85.8	82.18	88.70
<b>Form</b>					
Form 1	340	1196	81.8	75.63	86.66
Form 2	357	1240	83.2	76.07	88.59
Form 3	315	1097	82.9	78.34	86.63
Form 4	366	1025	87.6	80.36	92.41
Form 5	354	958	91.3	87.76	93.85
<b>Ethnicity</b>					
Malay	810	2540	87.3	83.78	90.08
Chinese	179	600	77.1	74.85	79.16
Indian	4	14	79.2	18.40	98.47
Bumiputera Sabah	474	1519	83.7	79.24	87.32
Bumiputera Sarawak	68	221	89.0	74.50	95.72
Others	197	623	85.8	76.43	91.87

- Prevalence with high RSE, not reported

**Table 3.6.11: Prevalence of handwashing most of the time or always after using the toilet in the past 30 days among adolescents in WP Labuan, 2022**

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
<b>WP LABUAN</b>	1789	5685	87.5	85.39	89.35
<b>Sex</b>					
Male	852	2680	84.7	80.27	88.27
Female	937	3004	90.2	88.40	91.70
<b>Form</b>					
Form 1	338	1189	81.3	76.97	84.91
Form 2	372	1283	86.2	78.83	91.25
Form 3	329	1144	86.4	81.45	90.23
Form 4	384	1078	92.1	87.69	94.98
Form 5	366	990	94.3	90.73	96.59
<b>Ethnicity</b>					
Malay	804	2516	86.4	82.66	89.48
Chinese	212	712	91.4	88.58	93.59
Indian	5	-	-	-	-
Bumiputera Sabah	489	1558	85.8	79.14	90.58
Bumiputera Sarawak	75	244	98.5	86.70	99.85
Others	204	638	87.8	80.38	92.70

- Prevalence with high RSE, not reported

**Table 3.6.12: Prevalence of handwashing using running water before eating at school in the past 30 days among adolescents in WP Labuan, 2022**

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
<b>WP LABUAN</b>	984	3131	48.2	42.17	54.37
<b>Sex</b>					
Male	417	1317	41.7	35.71	47.88
Female	567	1814	54.5	46.90	61.88
<b>Form</b>					
Form 1	141	501	34.2	26.05	43.50
Form 2	219	760	51.1	42.44	59.61
Form 3	177	617	46.8	38.19	55.53
Form 4	198	567	48.5	43.94	53.14
Form 5	249	686	65.3	46.49	80.31
<b>Ethnicity</b>					
Malay	431	1357	46.7	37.67	55.99
Chinese	89	290	37.3	28.99	46.43
Indian	2	-	-	-	-
Bumiputera Sabah	315	1009	55.6	49.63	61.36
Bumiputera Sarawak	45	146	59.0	39.96	75.61
Others	102	322	44.3	35.46	53.52

- Prevalence with high RSE, not reported

## 3.7 Mental Health Problems

### 3.7.1 Mental Health Problems

Contributors: Norhafizah Sahril, Muhammad Azri Adam Adnan, Kishwen Kanna Yoga Ratnam, Muhamad Khairul Nazrin Khalil, Mohamad Aznuddin Abd Razak, Sheikh Shafizal Sheikh Ilman, Chan Yee Mang, Mohd Shaiful Azlan Kassim, Sherina Mohd Sidik, Nurashikin Ibrahim, Raihan Khamal, Nor Rahidah Abdul Rahim, Noor Ani Ahmad.

#### 3.7.1.1 Introduction

According to the World Health Organization (WHO), one in every seven children and adolescents suffers from mental health problems, accounting for 13.0% of the global disease burden in this age group.<sup>1</sup> In Malaysia, the National Health and Morbidity Survey (NHMS) 2015 found that the prevalence of mental health problems was 34.7% among those aged 16 to 19, and 11.4% among those aged 10 to 15.<sup>2</sup> However, findings from the NHMS 2019 revealed that the prevalence of mental health problems had decreased to 9.5% among those aged 10 to 15 years old.<sup>3</sup> In particular, WHO reported that suicide is the fourth leading cause of death among 15-19 year-olds.<sup>1</sup> In 2017, 10.0% of secondary school adolescents reported suicidal ideation, according to the NHMS.<sup>4</sup>

#### 3.7.1.2 Objectives

- i. To identify the prevalence of loneliness in the past 12 months
- ii. To identify the prevalence of inability to sleep due to worry in the past 12 months
- iii. To identify the prevalence of suicidal ideation in the past 12 months
- iv. To identify the prevalence of suicidal plan in the past 12 months
- v. To identify the prevalence of suicidal attempt in the past 12 months
- vi. To identify the prevalence of not having close friends

#### 3.7.1.3 Variable definitions

- **Lonely “most of the time or always”:** Responded either “most of the time” or “always” for felt lonely during the past 12 months prior to the survey.
- **Unable to sleep “most of the time or always” due to worry:** Responded either “most of the time” or “always” for being worried about something that he/she could not sleep at night during the past 12 months prior to the survey.
- **Suicidal ideation:** ever seriously considered attempting suicide in the past 12 months prior to the survey.
- **Suicidal plan:** made a plan of attempted suicide in the past 12 months prior to the survey.
- **Suicidal attempt:** attempted suicide at least once in the past 12 months prior to the survey.
- **No close friend:** Do not have any close friend.

#### 3.7.1.4 Findings

Overall, 20.3% (95%CI: 17.89, 22.94) of adolescents in WP Labuan reported feeling lonely “most of the time or always” (**Table 3.7.1**). A total of 17.9% (95% CI: 13.60, 23.19) of adolescents reported being unable to sleep “most of the time or always” due to worry (**Table 3.7.2**). In the past 12 months prior to the survey, suicidal ideation, suicidal plan, and suicidal attempt, were reported 17.7% (95% CI: 13.74, 22.46), 14.0% (95% CI: 10.80, 17.93), and 13.0% (95% CI: 9.92, 16.95), respectively. (**Table 3.7.3**), (**Table 3.7.4**), (**Table 3.7.5**). The survey also observed that 3.6% (95% CI: 2.60, 4.85) of the adolescents had no close friends (**Table 3.7.6**).

#### 3.7.1.5 Discussion / Conclusion

The trend of suicidal ideation among secondary school students in WP Labuan (17.7%) was higher than GSHS2017 (12.3%) in 2017. In addition, this figure was also higher in comparison to national prevalence of 13.1%. Prevalence of suicidal plan in this survey (14.0%) was higher compared to the prevalence in GSHS 2017 (8.4%). This prevalence was also higher compared to the national prevalence of 10.0%. Prevalence of suicidal attempts in this survey (13.0%) was reported higher compared to the prevalence in than GSHS 2017 (6.9%). Moreover, this prevalence was higher compared to national prevalence (9.5%). In addition to these, less students (3.6%) in WP Labuan reported having no close friends as compared to previous GSHS 2017 (3.7%). This figure was also lower compared to national prevalence (4.2%).

#### 3.7.1.6 Recommendations

1. Enhanced the screening of at-risk adolescents by School Health Teams and referral for further management.
2. Intensify efforts to prevent suicide among student especially among high-risk group (attempt suicide).
3. Strengthen adolescents coping skills and resilience through interactive health promotion activities.
4. To introduce culturally competent programmes in school that upskill teachers and educate parents about risk of suicide among adolescents.
5. To improve the National school curriculum that teaches life skills such as effective coping strategies and develops mental resilience.
6. To review workplace policies with the aim of strengthening family ties such as the introduction of flexible working hours or the provision of options to work from home to increase quality time among parents and children.
7. To review school curriculum and teaching hours to optimize more time for physical activity and quality time for social and professional interaction among adolescents to improve adolescents’ life skills.

**Table: Mental Health Problems Trend in WP Labuan**

	NHMS 2017	NHMS 2022
Loneliness	9.8	20.3
Inability to sleep due to worry	9.4	17.9
Suicidal ideation	12.3	17.7
Suicidal plan	8.4	14.0
Suicidal attempt	6.9	13.0
Not having any close friend	3.7	3.6

**3.7.1.7 References**

1. WHO Fact Sheet. Adolescent mental health. <https://www.who.int/news-room/fact-sheets/detail/adolescent-mental-health>. Updated on 17 November 2021
2. Institute for Public Health (IPH). 2015. National Health and Morbidity Survey, NHMS 2015. Ministry of Health Malaysia
3. Institute for Public Health (IPH). 2019. National Health and Morbidity Survey, NHMS 2019. Ministry of Health Malaysia
4. Institute for Public Health (IPH). 2017. National Health and Morbidity Survey, NHMS 2017. Ministry of Health Malaysia



### 3.7.2 Depression

Contributors: Norhafizah Sahril, Muhamad Khairul Nazrin Khalil, Kishwen Kanna Yoga Ratnam, Muhammad Azri Adam Adnan, Mohamad Aznuddin Abd Razak, Sheikh Shafizal Sheikh Ilman, Chan Yee Mang, Mohd Shaiful Azlan Kassim, Sherina Mohd Sidik, Nurashikin Ibrahim, Raihan Khamal, Nor Rahidah Abdul Rahim, Noor Ani Ahmad.

#### 3.7.2.1 Introduction

Depression is a common mental health problem among adolescents worldwide. Depression can manifest as symptoms such as sadness, guilt, low self-esteem, a lack of happiness, and dissatisfaction with their surroundings.<sup>1</sup> Furthermore, depression can cause individual problems such as difficulty sleeping, loss of appetite, lack of energy, and easy despair, leading to suicidal ideation.<sup>2</sup> According to the World Health Organization (WHO), depression affects 1.1% of adolescents aged 10-14 years and 2.8% of those aged 15-19 years.<sup>3</sup> In Malaysia, the National Health and Morbidity Survey (NHMS) 2019 found that the prevalence of depression was 2.1% among those aged 15 to 19 years old.<sup>4</sup>

#### 3.7.2.2 Objectives

To determine the prevalence of depression among Malaysian adolescents.

#### 3.7.2.3 Variable definitions

**Depression:** A positive score was defined as a score of 10 and above for Patient Health Questionnaire (PHQ-9), and participants with these scores were categorized as having depression.

#### 3.7.2.4 Findings

Overall, 34.4% (95% CI: 28.89, 40.40) of WP Labuan adolescents reported depression. The prevalence of depression was significantly higher in female students 46.8% (95% CI: 40.59, 53.15) compared to males 21.3% (95% CI: 18.11, 24.91) (Table 3.7.7).

#### 3.7.2.5 Discussion / Conclusion

This survey indicated a higher prevalence of depression than research done by Normala et al. among 1800 Malaysian secondary school students aged 13 to 17 years old, which reported a 32.7% prevalence of depression among adolescents. Normala's study employed the same depression-measuring tool, the PHQ-9, but it was limited to 10 of 37 randomly chosen secondary schools in the Hulu Langat district area in the state of Selangor. Furthermore, the prevalence of depression in WP Labuan was higher compared to the national figure 26.9%.

### 3.7.2.6 Recommendations

1. Enhanced the screening of at-risk adolescents by School Health Teams and referral for further management.
2. Holistic intervention programmes targeted to adolescents at risk of depression.
3. Strengthen adolescents coping skills and resilience through interactive health promotion activities.
4. To introduce culturally competent programmes in school that upskill teachers and educate parents about discipline style and pro social parenting techniques.
5. To improve the National school curriculum that teaches life skills such as effective coping strategies and develops mental resilience.
6. To review workplace policies with the aim of strengthening family ties, such as the introduction of flexible working hours or the provision of options to work from home to increase quality time among parents and children.
7. To review school curriculum and teaching hours to optimize more time for physical activity and quality time for social and professional interaction among adolescents to improve adolescents' life skills.

### 3.7.2.7 References

1. Aquino JP, Londono A, Carvalho AF. An update on the epidemiology of major depressive disorder across cultures. In *Understanding depression 2018* (pp. 309-315). Springer, Singapore
2. Kaur J, Cheong SM, Mahadir Naidu B, Kaur G, Manickam MA, Mat Noor M, Ibrahim N, Rosman A. Prevalence and correlates of depression among adolescents in Malaysia. *Asia Pac J Public Health*. 2014 Sep;26(5\_suppl):53S-62S
3. WHO Fact Sheet. Adolescent mental health. <https://www.who.int/news-room/fact-sheets/detail/adolescent-mental-health>. Updated on 17 November 2021
4. Institute for Public Health (IPH). 2019. National Health and Morbidity Survey, NHMS 2019. Ministry of Health Malaysia

**Table 3.7.1: Prevalence of loneliness “most of the time or always” in the past 12 months among adolescents in WP Labuan, 2022**

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
<b>WP LABUAN</b>	415	1319	20.3	17.89	22.94
<b>Sex</b>					
Male	141	440	13.9	11.49	16.71
Female	274	879	26.4	23.05	29.99
<b>Form</b>					
Form 1	76	262	17.9	12.44	25.11
Form 2	101	343	23.0	17.97	29.00
Form 3	80	271	20.4	13.39	29.90
Form 4	87	246	21.1	17.81	24.71
Form 5	71	196	18.7	14.66	23.57
<b>Ethnicity</b>					
Malay	185	581	19.9	17.60	22.52
Chinese	38	126	16.1	12.35	20.79
Indian	0	-	-	-	-
Bumiputera Sabah	131	418	23.0	17.37	29.84
Bumiputera Sarawak	19	61	24.7	16.40	35.32
Others	42	133	18.3	12.74	25.69

- Prevalence with high RSE, not reported

**Table 3.7.2: Prevalence of inability to sleep “most of the time or always” due to worry in the past 12 months among adolescents in WP Labuan, 2022**

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
<b>WP LABUAN</b>	364	1163	17.9	13.60	23.19
<b>Sex</b>					
Male	112	347	11.0	9.36	12.78
Female	252	816	24.5	18.44	31.75
<b>Form</b>					
Form 1	64	222	15.2	9.25	23.95
Form 2	75	251	16.9	11.25	24.56
Form 3	82	284	21.5	12.57	34.17
Form 4	66	190	16.2	12.31	21.03
Form 5	77	215	20.5	13.43	29.98
<b>Ethnicity</b>					
Malay	159	504	17.3	12.95	22.78
Chinese	33	110	14.2	8.01	23.84
Indian	0	-	-	-	-
Bumiputera Sabah	127	405	22.3	14.78	32.15
Bumiputera Sarawak	15	48	19.3	10.90	31.92
Others	30	96	13.2	9.28	18.36

- Prevalence with high RSE, not reported

**Table 3.7.3: Prevalence of suicidal ideation in the past 12 months among adolescents in WP Labuan, 2022**

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
<b>WP LABUAN</b>	357	1149	17.7	13.74	22.46
<b>Sex</b>					
Male	75	233	7.4	5.46	9.89
Female	282	915	27.5	23.31	32.07
<b>Form</b>					
Form 1	78	272	18.6	12.41	26.84
Form 2	78	255	17.1	12.36	23.16
Form 3	77	260	19.7	13.76	27.31
Form 4	64	192	16.4	13.40	19.84
Form 5	60	171	16.2	8.58	28.60
<b>Ethnicity</b>					
Malay	129	406	14.0	9.45	20.14
Chinese	38	127	16.4	11.29	23.13
Indian	1	-	-	-	-
Bumiputera Sabah	121	394	21.7	14.09	31.96
Bumiputera Sarawak	20	64	25.9	15.19	40.57
Others	48	152	21.0	15.80	27.29

- Prevalence with high RSE, not reported

**Table 3.7.4: Prevalence of suicidal plan in the past 12 months among adolescents in WP Labuan, 2022**

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
<b>WP LABUAN</b>	278	908	14.0	10.80	17.93
<b>Sex</b>					
Male	59	195	6.2	3.92	9.58
Female	219	713	21.4	17.59	25.80
<b>Form</b>					
Form 1	68	238	16.3	10.33	24.73
Form 2	67	220	14.8	11.07	19.42
Form 3	56	193	14.6	8.77	23.36
Form 4	45	139	11.9	8.84	15.88
Form 5	42	118	11.2	6.32	19.22
<b>Ethnicity</b>					
Malay	108	344	11.8	8.23	16.75
Chinese	32	109	13.9	9.22	20.53
Indian	0	-	-	-	-
Bumiputera Sabah	81	268	14.8	8.88	23.56
Bumiputera Sarawak	16	52	20.8	10.63	36.72
Others	41	136	18.7	12.40	27.26

- Prevalence with high RSE, not reported

**Table 3.7.5: Prevalence of suicidal attempt "at least once" in the past 12 months among adolescents in WP Labuan, 2022**

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
<b>WP LABUAN</b>	259	847	13.0	9.92	16.95
<b>Sex</b>					
Male	61	205	6.5	4.79	8.71
Female	198	642	19.3	14.89	24.57
<b>Form</b>					
Form 1	61	213	14.5	11.13	18.77
Form 2	67	222	14.9	9.55	22.50
Form 3	55	186	14.1	9.05	21.22
Form 4	42	128	11.0	7.61	15.56
Form 5	34	98	9.3	4.62	17.90
<b>Ethnicity</b>					
Malay	98	317	10.9	7.76	15.06
Chinese	31	107	13.8	8.99	20.57
Indian	0	-	-	-	-
Bumiputera Sabah	84	274	15.1	10.09	22.00
Bumiputera Sarawak	12	-	-	-	-
Others	34	111	15.3	12.05	19.30

- Prevalence with high RSE, not reported

**Table 3.7.6: Prevalence of not having any close friends among adolescents in WP Labuan, 2022**

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
<b>WP LABUAN</b>	75	231	3.6	2.60	4.85
<b>Sex</b>					
Male	39	122	3.8	2.82	5.24
Female	36	109	3.3	1.82	5.85
<b>Form</b>					
Form 1	18	61	4.2	2.53	6.81
Form 2	16	54	3.6	2.30	5.73
Form 3	12	40	3.0	1.50	5.93
Form 4	15	-	-	-	-
Form 5	14	38	3.7	2.07	6.40
<b>Ethnicity</b>					
Malay	34	108	3.7	2.12	6.39
Chinese	8	26	3.3	1.94	5.51
Indian	0	-	-	-	-
Bumiputera Sabah	19	56	3.1	1.62	5.82
Bumiputera Sarawak	3	-	-	-	-
Others	11	33	4.6	2.44	8.34

- Prevalence with high RSE, not reported

**Table 3.7.7: Prevalence of depression among adolescents in WP Labuan, 2022**

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
<b>WP LABUAN</b>	695	2231	34.4	28.89	40.40
<b>Sex</b>					
Male	212	672	21.3	18.11	24.91
Female	483	1559	46.8	40.59	53.15
<b>Form</b>					
Form 1	130	456	31.3	25.49	37.68
Form 2	166	558	37.5	28.59	47.39
Form 3	147	500	37.8	25.68	51.59
Form 4	133	389	33.4	27.68	39.69
Form 5	119	327	31.3	23.89	39.71
<b>Ethnicity</b>					
Malay	297	942	32.4	25.23	40.60
Chinese	70	231	29.8	21.23	40.11
Indian	1	-	-	-	-
Bumiputera Sabah	227	731	40.4	33.59	47.50
Bumiputera Sarawak	25	81	32.7	18.04	51.86
Others	75	241	33.2	26.25	41.01

- Prevalence with high RSE, not reported

### 3.8 Physical Activity

Contributors: Muhammad Solihin Rezali, Affendi Isa, Siti Balkhis Shafie, Lim Kuang Kuay, Mohamad Aznuddin Abd Razak, Mohd Shaiful Azlan Kassim, Azli Baharudin@Shaharudin, Mohd Hairmansah Mohd Shah, Nor'Ain Ab Wahab, Norliza Shamsuddin, Nazirah Alias, Nurul Haniyah Rosslan, Hazizi Abu Saad, Mohd Azahadi Omar, Nur Hidayatun Fadhillah Mohd Nor

#### 3.8.1 Introduction

World Health Organization (WHO) defines physical activity as any bodily movement produced by skeletal muscles that require energy expenditure.<sup>1</sup> Based on WHO, at least 60 minutes per day of moderate-to-vigorous intensity physical activity were recommended for children and adolescents aged 5–17 years.<sup>1</sup> Sufficient physical activity has substantial health benefits for children and adolescents in terms of improving cardio-metabolic health, better musculoskeletal health, increased psychosocial well-being and academic performance.<sup>2</sup> Despite these established benefits, a substantial proportion of young people fail to meet physical activity guidelines. In addition, adolescents are also exposed to sedentary behaviours, as most of them spend greater time engaged in recreational activities, such as screen-based entertainment and digital communications.<sup>3</sup> Agenda National Malaysia Sihat (ANMS) and National Strategic Plan for Active Living (NASPAL) targeted to increase the adoption of healthy lifestyles among Malaysians which includes reducing the prevalence of physical inactivity among the general population, including adolescents by 10% within 10 years of implementation.<sup>4</sup> Thus, this study will provide more information regarding physical activity among adolescents in WP Labuan.

#### 3.8.2 Objectives

- i. To identify the prevalence of being physically active for a total of at least 60 minutes daily for five days or more in the past seven days among adolescents in WP Labuan
- ii. To identify the prevalence of active transportation or commuting among adolescents in WP Labuan
- iii. To identify the prevalence of sitting behavior among adolescents in WP Labuan

#### 3.8.3 Variable Definitions

- **Physically active:** physically active for at least 60 minutes per day, for a minimum of five days per week (sum of all the time spent in any kind of physical activity each day).
- **Active transportation/ commuting:** walking or riding a bicycle for at least three days a week to or from school.

- **Sitting behaviour:** Spending time sitting for 3 hours or more in a typical or usual day for leisure activities such as watching television, playing computer games, talking with friends, or surfing the internet.

#### 3.8.4 Findings

##### Physically active

The prevalence of being physically active was 22.4% (95% CI: 17.85, 27.67), among adolescents in WP Labuan. Males had significantly higher prevalence [27.4% (95% CI: 21.48, 34.35)] compared to females [17.6% (95% CI: 10.44, 28.03)]. (Table 3.8.1).

##### Active Transportation / Commuting

Overall, 17.8% (95% CI: 14.13, 22.09) adolescents reported active transportation to school. By state, the prevalence was highest in WP Putrajaya [46.9% (95% CI: 35.26, 58.87)], followed by WP Kuala Lumpur [36.6% (95% CI: 26.63, 47.85)] and Melaka [33.3% (95% CI: 23.65, 44.62)]. The prevalence was higher among males [20.2% (95% CI: 15.84, 25.35)] compared to females [15.5% (95% CI: 10.48, 22.25)]. (Table 3.8.2).

##### Sitting behaviour

A total of 68.9% (95% CI: 62.67, 74.49) had spent at least three hours in a typical or usual day engaging in sitting activities. The prevalence was higher in females [74.0% (95% CI: 64.63, 81.61)] compared to males. (Table 3.8.3).

#### 3.8.5 Discussion / Conclusion

The prevalence of being physically active and sitting behaviour among school adolescents in WP Labuan was higher than national findings (22.4% vs. 21.4%, 68.9% vs. 66.7%, respectively). In addition, the prevalence of active commuting was lower than the national finding (17.8% vs. 27.0%). Compared to previous NHMS findings, the prevalence of being physically active in the current study was lower than in the previous WP Labuan AHS 2017 (22.4% vs. 24.0%). The prevalence of sitting behaviour greatly increased from 2017 until the current survey (46.9% in 2017, and 68.9% in 2022).

#### 3.8.6 Recommendations

A comprehensive, integrated, intersectoral approach is required to increase the prevalence of physical activity among secondary school adolescents. Those initiatives and collaborative efforts jointly implemented across diverse ministries, agencies, private sectors, and civil service societies seem very effective, realizing that the social determinants of active living are beyond the health sectors. The recommendations are as below:

1. To explore more behavioral science and behavioral insights into physical inactivity and sedentary behavior among adolescents in WP Labuan to help us to design evidence-based health promotion and education initiatives with underlying effective 'nudging' techniques.

2. To examine the association between the frequency of use of online-related behaviors (time spent on social media, online communication, and e-games) with a sufficient level of physical activity and sedentary behavior in a more specific manner.
3. To include parents in the interventions and health promotion programs on physical activity among adolescents to encourage and support their children's participation in physical activity.

### 3.8.7 References

1. The World Health Organization. Physical Activity 2022 [Available from: <https://www.who.int/newsroom/fact-sheets/detail/physical-activity>]
2. Ekelund U, Luan Ja, Sherar LB, Esliger DW, Griew P, Cooper A, et al. Moderate to vigorous physical activity and sedentary time and cardiometabolic risk factors in children and adolescents. *JAMA*. 2012;307(7):704-12
3. Xu G, Sun N, Li L, Qi W, Li C, Zhou M, et al. Physical behaviors of 12-15 year-old adolescents in 54 low and middle-income countries: Results from the Global School-based Student Health Survey. *J. Glob. Health*. 2020;10(1)
4. Ministry of Health Malaysia. National Strategic Plan for Active Living (NASPAL) 2017-2025. Malaysia 2018

**Table 3.8.1: Prevalence of being physically active (at least 60 minutes daily) for a total of 5 days or more in the past 7 days in WP Labuan, 2022**

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
<b>WP LABUAN</b>	448	1454	22.4	17.85	27.67
<b>Sex</b>					
Male	20	869	27.4	21.48	34.35
Female	179	585	17.6	10.44	28.03
<b>Form</b>					
Form 1	73	260	17.8	12.49	24.61
Form 2	97	356	23.9	17.34	32.00
Form 3	92	326	24.6	15.96	35.91
Form 4	95	268	22.9	13.91	35.23
Form 5	91	245	23.3	19.17	28.04
<b>Ethnicity</b>					
Malay	187	588	20.2	15.26	26.21
Chinese	58	196	25.2	16.38	36.63
Indian	0	-	-	-	-
Bumiputera Sabah	131	428	23.6	17.37	31.18
Bumiputera Sarawak	24	83	33.5	20.41	49.79
Others	48	159	21.9	14.65	31.44

- Prevalence with high RSE, not reported

**Table 3.8.2: Prevalence of active commuting to school (walk or ride a bicycle to or from school for at least 3 days or more in the past 7 days) in WP Labuan, 2022**

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
<b>WP LABUAN</b>	366	1153	17.8	14.13	22.09
<b>Sex</b>					
Male	204	638	20.2	15.84	25.35
Female	162	515	15.5	10.48	22.25
<b>Form</b>					
Form 1	61	209	14.3	9.53	20.88
Form 2	75	262	17.7	12.79	23.85
Form 3	85	293	22.1	13.45	34.14
Form 4	80	218	18.6	13.31	25.48
Form 5	65	171	16.3	10.59	24.26
<b>Ethnicity</b>					
Malay	170	533	18.3	14.22	23.29
Chinese	30	97	12.5	8.96	17.18
Indian	1	-	-	-	-
Bumiputera Sabah	104	329	18.1	12.66	25.30
Bumiputera Sarawak	10	31	12.4	5.64	25.04
Others	51	159	22.0	11.20	38.59

- Prevalence with high RSE, not reported



**Table 3.8.3: Prevalence of spending at least 3 hours in sitting activities in WP Labuan, 2022**

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
<b>WP LABUAN</b>	1402	4472	68.9	62.67	74.49
<b>Sex</b>					
Male	636	2006	63.5	56.68	69.78
Female	766	2466	74.0	64.63	81.61
<b>Form</b>					
Form 1	245	866	59.2	51.99	65.97
Form 2	1	992	66.6	54.09	77.17
Form 3	280	972	73.4	62.88	81.84
Form 4	307	871	74.4	61.04	84.32
Form 5	284	771	73.8	65.27	80.88
<b>Ethnicity</b>					
Malay	619	1937	66.6	59.76	72.77
Chinese	176	592	76.0	67.80	82.63
Indian	2	-	-	-	-
Bumiputera Sabah	399	1283	70.7	62.35	77.84
Bumiputera Sarawak	54	176	70.8	49.93	85.46
Others	152	478	66.0	58.79	72.59

- Prevalence with high RSE, not reported

### 3.9 Protective Factors

Contributors: Nazirah Alias, LeeAnn Tan, Khaw Wan-Fei, S Maria Awaluddin, Eida Nurhadzira Muhammad, Filza Noor Asari, Mohd Amierul Fikri Mahmud, Faizul Akmal Abdul Rahim, Nur Hamizah Nasaruddin, Noor Syaqlah Shawaluddin, Mohd Farihan Md Yatim, Nik Rubiah Nik Abdul Rashid, Nik Daliana Nik Farid, Zamzaireen Zainal Abidin

#### 3.9.1 Introduction

Protective factors are individual or environmental characteristics or conditions that promote adolescent health and well-being<sup>1</sup>. The role of protective factors in adolescents is to improve the likelihood of positive health behaviours or outcomes (such as healthy diet, exercise, hygiene practices) and to reduce the negative impacts of risk factors (for example tobacco, alcohol and drug use, violence). Multiple protective factors at the school, peer and family levels can foster healthy behaviours and promote mental health<sup>2</sup>. At the school level, truancy is seen as an indicator that is monitored by lower prevalence, as truancy often acts as a precursor of many harmful behaviours. During adolescence, peer support and parental factors can be fundamental aspects of establishing positive health behaviours to prevent chronic diseases. In line with the strategies stated in the National Adolescent Health Policy, this study focuses on identifying protective factors at family, school, and peer levels that influence adolescent health and integrating these protective factors into health promotion among adolescents in WP Labuan.

#### 3.9.2 Objectives

- i. To determine the prevalence of truancy in the past 30 days among adolescents
- ii. To determine the prevalence of peer support in the past 30 days among adolescents
- iii. To determine the prevalence of parental or guardian supervision in the past 30 days among adolescents
- iv. To determine the prevalence of parental or guardian connectedness in the past 30 days among adolescents
- v. To determine the prevalence of parental or guardian bonding in the past 30 days among adolescents
- vi. To determine the prevalence of parental or guardian respect for privacy in the past 30 days among adolescents

#### 3.9.3 Variable definitions

- **Truancy:** Missed class or school without permission for at least one day in the past 30 days. (This variable is monitored with lower prevalence to define as protective factors).
- **Peer support:** Adolescents in their school were kind and helpful most of the time or always during the past 30 days.

- **Parental or guardian supervision:** Parents or guardians had always or most of the time, checked to see if their homework was done in the past 30 days.
- **Parental or guardian connectedness:** Parents or guardians had always or most of the time, understood their problems and worries in the past 30 days.
- **Parental or guardian bonding:** Parents or guardians had always or most of the time, really knew what they were doing with their free time in the past 30 days.
- **Parental or guardian respect for privacy:** Parents or guardians had never or rarely gone through their things without their approval in the past 30 days.

#### 3.9.4 Findings

##### Truancy

The prevalence of truancy in the past 30 days among adolescents was 29.8% (95% CI: 26.95, 32.87). It was higher in females (30.3%, 95% CI: 27.04, 33.79) compared to males (29.3%, 95% CI: 25.35, 33.64). Truancy was highest among Form 5 students with 36.5% (95% CI: 31.98, 41.24) (Table 3.9.1).

##### Having Peer Support

The prevalence of having peer support in the past 30 days among adolescents was 47.5% (95% CI: 42.21, 52.93). It was higher in females (57.5%, 95% CI: 51.83, 62.93) compared to males (37.1%, 95% CI: 34.68, 39.57). Having peer support was highest among Form 5 students with 57.4% (95% CI: 43.79, 69.95) (Table 3.9.2).

##### Having Parental or Guardian Supervision

The prevalence of having parental or guardian supervision in the past 30 days among adolescents was 8.1% (95% CI: 6.84, 9.47). It was higher in males (8.5%, 95% CI: 6.54, 11.07) compared to females (7.6%, 95% CI: 5.41, 10.58). Having parental or guardian connectedness was highest among Form 1 students with 10.9% (95% CI: 6.71, 17.18) (Table 3.9.3).

##### Having Parental or Guardian Connectedness

The prevalence of having parental or guardian connectedness in the past 30 days among adolescents was 18.9% (95% CI: 16.40, 21.59). It was higher in males (22.2%, 95% CI: 17.79, 27.29) compared to females (15.7%, 95% CI: 13.93, 17.64). Having parental or guardian connectedness was highest among Form 5 students with 22.6% (95% CI: 15.91, 31.11) (Table 3.9.4).

##### Having Parental or Guardian Bonding

The prevalence of having parental or guardian bonding in the past 30 days among adolescents was 27.2% (95% CI: 24.98, 29.52). It was higher in males (29.6%, 95% CI: 26.45, 33.05) compared to females (24.9%, 95% CI: 22.99, 26.84). Having parental or guardian bonding was highest among Form 5 students with 34.8% (95% CI: 29.95, 40.03) (Table 3.9.5).

**Having Parental or Guardian Respect for Privacy**

The prevalence of having parental or guardian respect for privacy in the past 30 days among adolescents was 81.9% (95% CI: 80.44, 83.24). It was higher in females (82.5%, 95% CI: 78.75, 85.78) compared to males (81.2%, 95% CI: 79.59, 82.69). Having parental or guardian respect for privacy was highest among Form 5 students with 87.7% (95% CI: 80.66, 92.37) (Table 3.9.6).

**3.9.5 Discussion / Conclusion**

Parental protective factors which were parent or guardian supervision, connectedness and bonding showed a decreasing trend. This is quite worrisome because parent/guardian-adolescent relation is a strong protective factor by providing a secure base for them especially in social support and might determine their children’s lives and behaviour during adolescence. Therefore, a comprehensive intervention policies or programmes must be further designed to address and to tackle this issue.

**3.9.6 Recommendations**

Development of interventions that strengthen the protective factors among school adolescents is important and more effective in reducing risk in order to improve the outcomes experienced by the adolescents. Among the interventions that can be implemented are:

1. Monitoring attendance closely by participation of schools, parent and local organizations through enforcement of mandatory attendance law allows identification of at risk and truancy behaviour among school adolescents.
2. Establishment of school programs that need parent’s supervision will help in improving the parenting skills especially in parental attachment.

**3.9.7 References**

1. Anthony, E. K., & Stone, S. I. (2010). Individual and contextual correlates of adolescent health and well-being. *Families in Society*, 91(3), 225–233. <https://doi.org/10.1606/1044-3894.3999>
2. Henson, M., Sabo, S., Trujillo, A., & Teufel-Shone, N. (2017). Identifying Protective Factors to Promote Health in American Indian and Alaska Native Adolescents: A Literature Review. *The journal of primary prevention*, 38(1-2), 5–26. <https://doi.org/10.1007/s10935-016-0455-2>

**Table: Protective Factors Trend in WP Labuan**

	NHMS 2012	NHMS 2017	NHMS 2022
Truancy	-	32.2	29.8
Having peer support	-	45.0	47.5
Having parental or guardian supervision	-	13.1	8.1
Having parental or guardian connectedness	-	31.5	18.9
Having parental or guardian bonding	-	38.0	27.2
Having parental or guardian respect for privacy	-	75.2	81.9

Table 3.9.1: Prevalence of truancy in the past 30 days among adolescents in WP Labuan, 2022

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
<b>WP LABUAN</b>	619	1938	29.8	26.95	32.87
<b>Sex</b>					
Male	300	928	29.3	25.35	33.64
Female	319	1010	30.3	27.04	33.79
<b>Form</b>					
Form 1	109	381	26.1	22.36	30.13
Form 2	118	406	27.3	20.92	34.74
Form 3	103	357	26.9	23.01	31.24
Form 4	149	411	35.1	29.22	41.41
Form 5	140	383	36.5	31.98	41.24
<b>Ethnicity</b>					
Malay	302	931	32.0	27.34	37.02
Chinese	59	198	25.4	17.58	35.21
Indian	0	-	-	-	-
Bumiputera Sabah	171	542	29.8	25.66	34.37
Bumiputera Sarawak	22	68	27.4	14.78	45.01
Others	65	199	27.5	21.86	33.87

- Prevalence with high RSE, not reported

Table 3.9.2: Prevalence of having peer support in the past 30 days among adolescents in WP Labuan, 2022

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
<b>WP LABUAN</b>	972	3089	47.5	42.21	52.93
<b>Sex</b>					
Male	376	1174	37.1	34.68	39.57
Female	596	1915	57.5	51.83	62.93
<b>Form</b>					
Form 1	162	570	39.0	32.90	45.38
Form 2	206	705	47.3	35.69	59.28
Form 3	182	626	47.3	40.20	54.47
Form 4	203	586	50.0	38.06	61.98
Form 5	219	603	57.4	43.79	69.95
<b>Ethnicity</b>					
Malay	435	1358	46.7	43.49	49.84
Chinese	100	329	42.3	24.67	62.08
Indian	1	-	-	-	-
Bumiputera Sabah	287	921	50.7	41.84	59.57
Bumiputera Sarawak	37	122	49.3	30.02	68.73
Others	112	355	48.9	42.29	55.64

- Prevalence with high RSE, not reported

**Table 3.9.3: Prevalence of having parental or guardian supervision in the past 30 days among adolescents in WP Labuan, 2022**

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
<b>WP LABUAN</b>	159	523	8.1	6.84	9.47
<b>Sex</b>					
Male	82	270	8.5	6.54	11.07
Female	77	253	7.6	5.41	10.58
<b>Form</b>					
Form 1	45	159	10.9	6.71	17.18
Form 2	42	147	9.9	6.54	14.72
Form 3	25	-	-	-	-
Form 4	24	65	5.5	2.78	10.76
Form 5	23	61	5.8	3.14	10.59
<b>Ethnicity</b>					
Malay	70	227	7.8	6.06	9.97
Chinese	18	63	8.1	4.74	13.51
Indian	0	-	-	-	-
Bumiputera Sabah	43	141	7.8	5.42	11.06
Bumiputera Sarawak	6	-	-	-	-
Others	22	71	9.8	5.66	16.56

- Prevalence with high RSE, not reported

**Table 3.9.4: Prevalence of having parental or guardian connectedness in the past 30 days among adolescents in WP Labuan, 2022**

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
<b>WP LABUAN</b>	386	1224	18.9	16.40	21.59
<b>Sex</b>					
Male	222	702	22.2	17.79	27.29
Female	164	522	15.7	13.93	17.64
<b>Form</b>					
Form 1	78	276	18.8	17.51	20.24
Form 2	75	267	17.9	12.81	24.52
Form 3	65	231	17.5	12.10	24.56
Form 4	79	214	18.2	12.27	26.25
Form 5	89	237	22.6	15.91	31.11
<b>Ethnicity</b>					
Malay	180	566	19.5	14.93	24.95
Chinese	42	142	18.2	12.80	25.22
Indian	3	-	-	-	-
Bumiputera Sabah	110	351	19.4	15.52	23.88
Bumiputera Sarawak	11	-	-	-	-
Others	40	121	16.7	11.30	23.86

- Prevalence with high RSE, not reported

Table 3.9.5: Prevalence of having parental or guardian bonding in the past 30 days among adolescents in WP Labuan, 2022

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
<b>WP LABUAN</b>	559	1767	27.2	24.98	29.52
<b>Sex</b>					
Male	300	938	29.6	26.45	33.05
Female	259	828	24.9	22.99	26.84
<b>Form</b>					
Form 1	108	380	26.0	22.17	30.17
Form 2	105	376	25.3	18.02	34.22
Form 3	97	339	25.6	21.86	29.74
Form 4	113	306	26.1	21.59	31.21
Form 5	136	366	34.8	29.95	40.03
<b>Ethnicity</b>					
Malay	263	822	28.2	23.31	33.72
Chinese	57	187	24.0	18.92	30.04
Indian	2	-	-	-	-
Bumiputera Sabah	167	533	29.4	25.33	33.76
Bumiputera Sarawak	20	63	25.4	16.32	37.25
Others	50	155	21.3	15.48	28.56

- Prevalence with high RSE, not reported

Table 3.9.6: Prevalence of having parental or guardian respect for privacy in the past 30 days among adolescents in WP Labuan, 2022

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
<b>WP LABUAN</b>	1671	5320	81.9	80.44	83.24
<b>Sex</b>					
Male	811	2570	81.2	79.59	82.69
Female	860	2750	82.5	78.75	85.78
<b>Form</b>					
Form 1	319	1116	76.3	68.23	82.82
Form 2	346	1201	80.6	77.48	83.39
Form 3	318	1107	83.7	78.50	87.76
Form 4	350	975	83.3	77.34	87.95
Form 5	338	920	87.7	80.66	92.37
<b>Ethnicity</b>					
Malay	758	2364	81.2	79.61	82.70
Chinese	197	666	85.5	81.12	89.08
Indian	4	14	77.8	17.19	98.34
Bumiputera Sabah	465	1489	82.0	77.13	86.03
Bumiputera Sarawak	57	189	76.0	64.21	84.86
Others	190	599	82.5	74.27	88.47

### 3.10 Sexual Behaviours

Contributors: Noor Aliza Lodz, Amal Shamsudin, Chong Zhuo Lin, Fatin Athira Tahir, Mazliza Ramly, Maznieda Mahjom, Nik Adilah Shahein, S Maria Awaluddin, Anita Suleiman, Nik Rubiah Nik Abdul Rashid

#### 3.10.1 Introduction

Adolescent sexual behaviour contributes to various sexual and reproductive health issues. The Global Summary HIV Epidemic Report, there were 150,000 adolescents aged 10-19 that were newly infected with HIV while 1,750,000 adolescents were already living with HIV<sup>1</sup>. It was also reported that the highest rate of sexual transmitted illness (STI) worldwide is among young people aged 15 to 24 years<sup>2</sup>. In Malaysia, the incidence of HIV reported among adolescents aged 13 to 19 was 2.4 per 100,000 population in 2021 and this trend has been steadily increasing in the past 10 years<sup>2</sup>. The WHO reported that 50% of young unmarried girls aged 15 to 19 years in low- and middle-income countries had an unintended pregnancy in 2019<sup>3</sup>. Globally, there were 41 births per 1000 girls aged 15-19 years in 2020 and 14% of maternal deaths. Good knowledge of HIV would help in reducing the transmission as shown in a study that a person with inadequate knowledge of HIV is more vulnerable to acquire the infection and may spread the disease throughout the population<sup>4</sup>. Good knowledge on HIV transmission was also associated with intention to engage in low-risk sexual behaviour<sup>5</sup>. Therefore, this study also aims to assess the knowledge of HIV transmission based on United Nation General Assembly Special Session (UNGASS) indicators among the school adolescents to determine the prevalence of HIV knowledge among them.

#### 3.10.2 Objectives

To determine:

- i. the prevalence of ever having sexual intercourse among adolescents in Malaysia
- ii. the prevalence of current sexual intercourse in the past 30 days among adolescents in Malaysia
- iii. the percentage of first sexual experience before the age 14 years among those who ever had sex
- iv. the percentage of having at least two sexual partners among those who ever had sex
- v. the percentage of condom usage during the last sexual intercourse among those who ever had sex
- vi. the percentage of "other birth control methods" usage during the last sexual intercourse among those who ever had sex
- vii. the prevalence of adequate HIV knowledge among adolescents in Malaysia
- viii. the percentage of correct responses in each of UNGASS indicators among adolescents in Malaysia

#### 3.10.3 Variable Definitions

- **Sexual intercourse:** sexual acts of penile penetration into the vagina or anus.
- **Risky sexual behaviour:** behaviours such as early sex debut, multiple sex partners and unprotected sex that could lead to health problems.
- **Other birth control methods:** pregnancy prevention methods other than barrier methods (condom usage) including withdrawal, birth control pills or any other non-barrier methods.
- **Ever had sex:** any positive answer for first sexual intercourse.
- **Current sexual intercourse:** sexual intercourse in the past 30 days.
- **Adequate HIV Knowledge:** provided correct responses to all five items of UNGASS indicators/questions.

#### 3.10.4 Findings

Prevalence of ever had sex among adolescents in WP Labuan was 7.1% (95% CI: 5.53, 9.16), male adolescents showed significantly higher prevalence 9.1% (95% CI: 6.80, 11.94) compared to females, 5.3% (95% CI: 4.02, 7.01) (**Table 3.10.1**). Prevalence of current sexual intercourse among adolescents in WP Labuan was 4.5% (95% CI: 3.44, 5.85), male adolescents had higher prevalence of currently having sexual intercourse which was 5.3% (95% CI: 3.61, 7.56) compared to female; 3.8% (95% CI: 2.78, 5.13) (**Table 3.10.2**). Of those who ever had sex, 39.1% had sex before the age of 14. It was noted that 12.7% of them used other birth control methods and 9.1% used condom during their last sexual intercourse. While 8.5% those who had at least two sexual partners (**Table 3.10.3**).

Overall, the prevalence of adequate HIV knowledge among adolescents was 1.7% (95% CI: 1.12, 2.70). Male adolescents prevalence with 2.0% (95% CI: 0.98, 4.00) of HIV knowledge (**Table 3.10.4**). The percentage of correct responses by each item was highest for question "Can a healthy-looking person have HIV?" with 25.7%. "Can a person get HIV from mosquito bites?" with percentage 20.7%. Followed by question "Can a person reduce the risk of getting HIV using a condom every time they have sex?" the percentage was 17.5% and 17.2% for "Can a person get HIV by sharing food with someone who is infected?". While the least 13.5% for "Can the risk of HIV transmission be reduced by having sex with only one uninfected partner who has no other partner?" (**Table 3.10.5**)

### 3.10.5 Discussion / Conclusion

This study found that majority of sexually active adolescents are engaging in risky sexual behaviour, i.e., sex debut before age 14 years, having multiple sexual partners and unprotected sex. While from the UNGASS indicators, adequate knowledge on HIV transmission among adolescents are still low.

### 3.10.6 Recommendations

1. To strengthen sexual and reproductive health education to be more effective and comprehensive in empowering adolescents with appropriate knowledge, attitude, and skills.
2. To enhance the promotion of various existing sexual and reproductive health modules designed to guide and assist parents / guardians / caregivers to talk about sexuality at home and institutions.
3. To improve on parenting skills and effective communication in sexual and reproductive health related matters.
4. To conduct more studies especially qualitative studies in exploring the determinants of risky sexual behaviours among adolescents.
5. To reactivate the Healthy Programme Without AIDS for Adolescents (PROSTAR) to increase HIV/STI awareness and knowledge.
6. To utilise creative and innovative approaches through social media, peer educator programmes, public-private-NGO (triparty) partnerships and etc.
7. To utilise creative and innovative approaches through social media, peer educator programmes, public-private-NGO (triparty) partnerships and etc.

### 3.10.7 References

1. HIV and AIDS in adolescents. Unicef Data. 2021 <https://data.unicef.org/topic/hiv/aids/#:~:text=Globally%2C%20adolescents%2010%2D19%20years,of%20all%20AIDS%2Drelated%20deaths>
2. WHO fact sheet: Adolescent pregnancy. 2022. <https://www.who.int/news-room/fact-sheets/detail/adolescent-pregnancy>
3. Lindberg LD, Firestein L, Beavin C. Trends in U.S. adolescent sexual behavior and contraceptive use, 2006-2019. *Contracept X*. 2021 Apr 8; 3:100064
4. T Carine Ronsmans, Wendy J Graham, on behalf of The Lancet Maternal Survival Series steering group, 2006. Maternal mortality: who, when, where, and why. *The Lancet's Maternal Survival and Women Deliver Series 2006/2007: 2005 World Health Report*
5. Satterwhite CL, Torrone E, Meites E, Dunne EF, Mahajan R, Ocfemia MC, et al. Sexually transmitted infections among US women and men: prevalence and incidence estimates, 2008. *Sex Transm Dis*. 2013;40(3):187-93



**Table 3.10.1: Prevalence of ever had sexual intercourse among adolescents in WP Labuan, 2022**

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
<b>WP LABUAN</b>	147	464	7.1	5.53	9.16
<b>Sex</b>					
Male	92	286	9.1	6.80	11.94
Female	55	177	5.3	4.02	7.01
<b>Form</b>					
Form 1	45	156	10.7	7.27	15.37
Form 2	28	101	6.8	5.10	9.05
Form 3	15	51	3.8	1.90	7.54
Form 4	35	92	7.8	4.59	13.02
Form 5	24	64	6.1	3.61	10.10
<b>Ethnicity</b>					
Malay	77	236	8.1	5.92	11.03
Chinese	10	-	-	-	-
Indian	0	-	-	-	-
Bumiputera Sabah	36	119	6.6	4.66	9.22
Bumiputera Sarawak	6	-	-	-	-
Others	18	-	-	-	-

- Prevalence with high RSE, not reported

**Table 3.10.2: Prevalence of current sexual intercourse in the past 30 days among adolescents in WP Labuan, 2022**

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
<b>WP LABUAN</b>	94	292	4.5	3.44	5.85
<b>Sex</b>					
Male	55	166	5.3	3.61	7.56
Female	39	126	3.8	2.78	5.13
<b>Form</b>					
Form 1	26	89	6.1	3.33	10.95
Form 2	18	64	4.3	2.72	6.66
Form 3	8	-	-	-	-
Form 4	23	61	5.3	3.05	8.89
Form 5	19	51	4.9	2.72	8.54
<b>Ethnicity</b>					
Malay	47	144	4.9	3.35	7.22
Chinese	5	-	-	-	-
Indian	0	-	-	-	-
Bumiputera Sabah	23	74	4.1	2.49	6.66
Bumiputera Sarawak	5	-	-	-	-
Others	14	-	-	-	-

- Prevalence with high RSE, not reported

**Table 3.10.3: Proportion of sexual practices among those who ever had sex among adolescents in WP Labuan, 2022**

Sexual Practices	Unweighted count	Percentage (%)
Percentage of first sex before the age 14 years	55	39.1
Percentage of having at least two sexual partners	13	8.5
Percentage of reported condom use during last sexual intercourse	12	9.1
Percentage of reported using other birth control method during last sexual intercourse	17	12.7

**Table 3.10.4: Prevalence of adequate HIV knowledge among adolescents by Sociodemographic Characteristics in WP Labuan, 2022**

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
<b>WP LABUAN</b>	37	113	1.7	1.12	2.70
<b>Sex</b>					
Male	22	63	2.0	0.98	4.00
Female	15	-	-	-	-
<b>Form</b>					
Form 1	3	-	-	-	-
Form 2	4	-	-	-	-
Form 3	5	-	-	-	-
Form 4	13	-	-	-	-
Form 5	12	-	-	-	-
<b>Ethnicity</b>					
Malay	9	25	0.9	0.52	1.47
Chinese	7	23	2.9	1.74	4.80
Indian	0	-	-	-	-
Bumiputera Sabah	14	41	2.3	1.28	4.02
Bumiputera Sarawak	0	-	-	-	-
Others	7	-	-	-	-

- Prevalence with high RSE, not reported

**Table 3.10.5: Percentage of Correct Responses by item of UNGASS Indicator among adolescents in WP Labuan, 2022**

Sexual Practices	Unweighted count	Percentage (%)
Can the risk of HIV transmission be reduced by having sex with only one uninfected partner who has no other partners?	281	13.5
Can a person reduce the risk of getting HIV using a condom every time they have sex?	359	17.5
Can a healthy-looking person have HIV?	540	25.7
Can a person get HIV from mosquito bites?	428	20.7
Can a person get HIV by sharing food with someone who is infected?	355	17.2

### 3.11 Tobacco Use

Contributors: Mohd Ruhaizie Riyadzi, Muhammad Fadhli Mohd Yusoff, Hamizatul Akmal Abd Hamid, Ummi Nadiah Yusoff, Nizam Baharom, Thamil Arasu Saminathan, Tania Galye Robert Lourdes, Halizah Mat Rifin, Lim Kuang Hock, Norliana Ismail, Muhammad Hairul Nizam Abd Hamid and Noraryana Hassan

#### 3.11.1 Introduction

Tobacco use including cigarette and e-cig/vape is predominantly an issue for male adolescents. Malaysia is committed to achieve smoke free generation by 2040<sup>1</sup>. Various anti-tobacco programs for youths have been established especially at school level. Continuous surveillance of tobacco use among adolescents is essential in monitoring the progress of tobacco control programs in Malaysia generally and at state level of WP Labuan, specifically.

#### 3.11.2 Objectives

##### General objective:

To determine the use of tobacco among adolescents in WP Labuan.

##### Specific objectives:

- i. To identify the prevalence of the current use of any tobacco product adolescents in WP Labuan
- ii. To identify the prevalence of the current e-cig/vape use among adolescent in WP Labuan
- iii. To determine the latest source of cigarette obtained among adolescent cigarette smokers in WP Labuan
- iv. To determine the latest source of e-cig/vape obtained among adolescent e-cig/vape users in WP Labuan
- v. To determine the prevalence of exposure to second-hand smoke among adolescent in WP Labuan
- vi. To determine the prevalence of exposure to tobacco products advertisement or promotion in the point of sales among adolescents in WP Labuan

#### 3.11.3 Variable Definitions

- **Current any tobacco use** - the use any of the following tobacco product during the last 30 days: manufactured cigarette, traditional hand rolled cigarettes, roll-your-own cigarettes with cigarette papers, cigar/cigarillos, tobacco pipe (pipe smoking), shisha/hookah, electronic cigarette/vape, heated tobacco product, snuff or chewed tobacco
- **Current tobacco smoker or current smoker** - the use of any of the following tobacco products during the last 30 days: manufactured cigarette, traditional hand rolled cigarettes, roll-your-own cigarettes with cigarette papers, cigar/cigarillos, tobacco pipe (pipe smoking) or shisha/hookah

- **Current cigarette smoker** - the use of any of the following tobacco products during the last 30 days: manufactured cigarettes, traditional hand rolled cigarettes, roll-your-own cigarettes with cigarette papers or cigar/cigarillos
- **Current e-cig/vape user** - the use of e-cig/vape during the last 30 days

#### 3.11.4 Findings

In WP Labuan, the prevalence of current use of any tobacco products was 22.2% (95%CI: 13.19, 34.91), which was above the national prevalence [18.5% (95%CI: 17.09, 19.92)]. The prevalence of males was two and half times higher [32.0% (95%CI: 16.64, 52.66)] as compared to females [12.9% (95%CI: 8.30, 19.45)] (**Table 3.11.1**). The prevalence of the e-cig/vape user among adolescents in WP Labuan was 19.2 (95%CI: 10.51, 32.32) with the prevalence of males was almost three times higher compared to females [28.3% (95%CI: 13.52, 49.88) vs. 10.5% (95%CI: 6.13, 17.33)] (**Table 3.11.2**).

Among those who smoked cigarettes, most of the cigarettes obtained by buying them from static premises (34.2%), by getting them from friends (29.0%) and by buying them from non-static premises (13.4%) (**Table 3.11.3**). Most of the e-cig/vape were obtained by getting them from friends (35.9%) and by buying them from specific e-cig/vape shops (35.8%) (**Table 3.11.4**). Nearly half of the adolescents reported they have parent or guardian who smoked or used any type of tobacco products [43.5% (95%CI: 38.06, 49.00)] and reported they were exposed to second-hand smoke when someone else smoking nearby in their presence within the past 7 days [42.6% (95%CI: 32.43, 53.39)], with more than a quarter [26.7% (95%CI: 23.33, 30.45)] reported they have e-cig/vape used parent or guardians (**Table 3.11.5**). Nearly one-fifth of the adolescents claimed they were exposed to the tobacco products point-of-sale advertising and promotion for the past 30 days [17.3% (95%CI: 15.20, 19.72)] (**Table 3.11.6**).

#### 3.11.5 Discussion / Conclusion

The prevalence of tobacco use among adolescents in WP Labuan just has some narrow difference as compared with the prevalence in Malaysia. There was a significant increase in the prevalence of e-cig/vape use among adolescents, as compared with findings in 2017<sup>2</sup>. This indicates a switch in the preference of nicotine delivery among adolescents in Malaysia, as well as in WP Labuan within the past five years. Various factors could have contributed to these recent findings. Special concern should also be given to female adolescents as the prevalence of e-cig/vape users has doubled since 2017<sup>2</sup>.

#### 3.11.6 Recommendation

Tobacco use, which includes vaping, is a major harmful determinant for human health. It is worrying that the current anti-tobacco programs seem to have not diminished

Malaysian adolescents' interest in vaping. A smokefree generation requires participation and dedication from all sectors, including family institutions, education sector, politicians, government and non-governmental organizations. Moving forward, it is high time for all sectors to come together and agree on banning tobacco use among future Malaysian generations. Strengthening the current law and taking legal action are vital in controlling the accessibility of tobacco products, especially e-cig/vape by adolescents. For those who have developed a nicotine addiction, the visibility of quit smoking services should be increased more aggressively to attract more adolescents to seek help. All screening, prevention, and intervention programs among adolescents must be strengthened and delivered in synergy by all governmental and nongovernmental agencies.

### 3.11.7 References

1. Tobacco Control Sector and FCTC Secretariat, 2021. National Strategic Plan for The Control of Tobacco & Smoking Products 2021-2030. Ministry of Health Malaysia
2. Institute for Public Health (IPH). 2017. National Health and Morbidity Survey, NHMS 2017. Ministry of Health Malaysia

**Table 3.11.1: Prevalence of current any tobacco use among adolescents in WP Labuan, 2022**

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
<b>WP LABUAN</b>	477	1443	22.2	13.19	34.91
<b>Sex</b>					
Male	342	1014	32.0	16.64	52.66
Female	135	429	12.9	8.30	19.45
<b>Form</b>					
Form 1	74	256	17.5	11.89	25.06
Form 2	72	263	17.6	9.96	29.32
Form 3	67	235	17.7	10.07	29.28
Form 4	139	-	-	-	-
Form 5	125	-	-	-	-
<b>Ethnicity</b>					
Malay	255	768	26.4	14.94	42.25
Chinese	20	-	-	-	-
Indian	0	-	-	-	-
Bumiputera Sabah	120	-	-	-	-
Bumiputera Sarawak	19	-	-	-	-
Others	63	194	26.7	16.85	39.65

- Prevalence with high RSE, not reported

**Table 3.11.2: Prevalence of current e-cig/vape use among adolescents in WP Labuan, 2022**

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
<b>WP LABUAN</b>	419	1244	19.2	10.51	32.32
<b>Sex</b>					
Male	309	895	28.3	13.52	49.88
Female	110	349	10.5	6.13	17.33
<b>Form</b>					
Form 1	57	197	13.5	8.50	20.68
Form 2	61	221	14.8	8.19	25.31
Form 3	52	179	13.5	6.70	25.30
Form 4	134	-	-	-	-
Form 5	115	-	-	-	-
<b>Ethnicity</b>					
Malay	225	665	22.8	11.56	40.12
Chinese	14	-	-	-	-
Indian	0	-	-	-	-
Bumiputera Sabah	110	-	-	-	-
Bumiputera Sarawak	18	-	-	-	-
Others	52	156	21.4	12.12	35.06

- Prevalence with high RSE, not reported

**Table 3.11.3: Proportion of source of cigarette obtaining during the last time smoking in the past 30 days among adolescents in WP Labuan, 2022**

Socio-demographic characteristics	Unweighted count	Percentage (%)
Bought from static premises	72	34.2
Bought from non-static premises	26	13.4
Food establishment	4	-
Bought online	9	-
Get from friends	62	29.0
Get from family members	13	-
Got some other ways	19	-

- Prevalence with high RSE, not reported

**Table 3.11.4: Proportion of source of e-cigarette/vape obtaining during the last time using e-cig/vape in the past 30 days among adolescents in WP Labuan, 2022**

Socio-demographic characteristics	Unweighted count	Percentage (%)
Bought from static premises	130	35.8
Bought from non-static premises	11	-
Food establishment	15	-
Bought online	13	-
Get from friends	131	35.9
Get from family members	31	-
Got some other ways	22	6.2

- Prevalence with high RSE, not reported

**Table 3.11.5: Prevalence of exposure to second hand smoke among adolescents in WP Labuan, 2022**

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
Having parent or guardian who smoked or used any tobacco products	819	2617	43.5	38.06	49.00
Having e-cigarette/vape use parent or guardian	506	1632	26.7	23.33	30.45
Exposure to second hand smoke in the past 7 days	896	2766	42.6	32.43	53.39

**Table 3.11.6: Prevalence of currently see or notice any tobacco products advertising or promotion in the past 30 days among adolescents in WP Labuan, 2022**

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
Currently see or notice any tobacco product advertising or promotion in the point of sales in the past 30 days	362	1126	17.3	15.20	19.72

### 3.12 Violence and Unintentional Injury

Contributors: Hamizatul Akmal Abd Hamid, Tan Lee Ann, Nor Rahidah Abd Rahim, Noor Raihan Khamal, Mohd Hazrin Bin Hasim@Hashim, Nur Faraeein Zainal Abidin, Muhammad Hanafi Bakri, Noor Suraya Muhamad, Shubash Shander Ganapathy

#### 3.12.1 Introduction

Malaysia supports the mandate under resolutions WHA67.15 (2014) and WHA69.5 (2016) on implementing the WHO global plan of action to strengthen the role of the health system within a national multisectoral response to address interpersonal violence in particular against women and girls, and against children. Global school-based health surveys have shown that up to 42% of adolescent boys and 37% of adolescent girls were exposed to bullying<sup>1</sup>. Due to lockdowns caused by the ongoing COVID-19 pandemic, adolescents may be subjected to mistreatment and violence when they are forced to remain at home with their aggressors<sup>2</sup>. In addition, cyberbullying is another issue of concern that is closely related to adolescents' mental health and development<sup>3</sup>.

#### 3.12.2 Objectives

To describe the prevalence of:

- i. Having been physically attacked at least once in the past 12 months
- ii. Involvement in a physical fight at least once in the past 12 months
- iii. Having had a serious injury at least once in the past 12 months
- iv. Physical abuse at home at least once in the past 30 days
- v. Verbal abuse at home at least once in the past 30 days
- vi. Having been bullied at least once in the past 30 days
- vii. Involvement in the perpetration of cyberbullying a few times within a year or more

#### 3.12.3 Variable Definitions

- **Physical attack:** when one or more persons hurt another person with or without a weapon such as sticks or knives in the past 12 months. It is NOT a physical attack when two individuals or adolescents of about the same strength or power choose to fight each other.
- **Physical fight:** when two individuals or adolescents of about the same strength or power choose to fight each other in the past 12 months.
- **Unintentional injury:** a serious injury which makes the student miss at least one full day of usual activity (such as school, sports or a job) OR requires treatment by doctor or medical personnel in the past 12 months.

- **Physical abuse at home:** when someone is hit so hard that it left a mark OR caused an injury in the past 30 days.
- **Verbal abuse at home:** when someone has had hurtful or insulting things said to them in the past 30 days.
- **Bullying:** when a student or group of adolescents say or do bad and unpleasant things to another student, such as teasing a lot in an unpleasant way or leaving out things on purpose in the past 30 days. It is NOT bullying when two adolescents of about the same strength or power argue or fight or when teasing is done in a friendly and fun way.
- **Cyberbullying (perpetrator):** bullying or harassment through the internet, cell phones, or other electronic devices (ie, sending insulting messages, posting digitally altered photos, engaging in online fighting, making aggressive comments, sharing someone's embarrassing information, or sending messages that include threats of harm through e-mail, instant messaging, in a chat room, on a website, or sent to a cell phone).

#### 3.12.4 Findings

The prevalence of adolescents who had been physically attacked in the past 12 months was 17.4% (95% CI: 15.07, 20.02) and this was similar in both in female [17.4% (95% CI: 14.42, 20.86)] and male [17.4% (95% CI: 13.86, 21.63)]. (**Table 3.12.1**). Overall, 20.5% (95% CI: 18.89, 22.29) adolescents claimed to have been involved in a physical fight, which was higher in male [22.7% (95% CI: 19.02, 26.89)] compared to female [18.5% (95% CI: 15.49, 21.88)]. (**Table 3.12.1**).

The prevalence of adolescents who had a serious injury in the past 12 months was 20.2% (95% CI: 18.57, 21.87). Male adolescents showed a higher prevalence [21.8% (95% CI: 18.63, 25.24)] compared to female [18.7% (95% CI: 16.71, 20.79)]. (**Table 3.12.2**). However, among those who had been seriously injured, the two most common causes of serious injury were falls [7% (95% CI: 5.80, 8.41)] and something fell or hit him/her [1.1% (95% CI: 0.79, 1.54)]. (**Table 3.12.3**).

The prevalence of adolescents reported had experienced physical abuse at home was 9.9% (95% CI: 7.84, 12.49) which was higher among female at 11.5% (95% CI: 7.66, 16.86) compared to male at 8.3% (95% CI: 5.32, 12.70)]. (**Table 3.12.4**). Overall, 47% (95% CI: 40.71, 53.35) adolescents reported being abused verbally at home and it was significantly higher among female [62% (95% CI: 58.12, 65.77)] compared to male [31.2% (95% CI: 28.21, 34.26)]. (**Table 3.12.4**).

In terms of bullying, 9.9% (95% CI: 8.28, 11.87) adolescents reported having been bullied. This was higher among female [11.2% (95% CI: 8.48, 14.60)] compared to male [8.6% (95% CI: 6.08, 12.08)]. (**Table 3.12.5**). The most

common form of bullying was, 'Making fun of how my body or face looks' [2% (95% CI: 1.56, 2.51)]. (Table 3.12.6).

With regards to involvement in cyberbullying activities from the perspective of the perpetrator, 21.2% (95% CI: 17.89, 24.89) of adolescents reported that they had been involved in cyberbullying activities a few times within the past year or more. Male adolescents showed a significantly higher prevalence [25.9% (95% CI: 21.40, 30.88)] compared to female [16.7% (95% CI: 14.18, 19.62)]. (Table 3.12.7). The two most common forms of adolescents' involvement in cyberbullying activities were 'Ever made rude comments to anyone online' [12.9% (95% CI: 10.06, 16.47)] and 'Ever spread rumours about someone online' [7.4% (95% CI: 5.79, 9.41)]. (Table 3.12.8).

### 3.12.5 Discussion / Conclusion

Pandemic situations reduced the prevalence of violence significantly through restricted movement, reduced social exposure, and reduced exposure to physical violence. The prevalence of perpetration of cyberbullying found in this survey is consistent with only a few countries in the world. Despite the reduction in all domains, further strategic steps should be taken to improve the outcome of the survey.

### 3.12.6 Recommendations

In the previous two surveys, recommendations touched on identifying the risk factors that contributed to the problems. The recommendations in this survey are more focused on dealing with abuse, bullying, cyberbullying, and falling. Approaches should be comprehensive with the involvement of relevant agencies.

- i. Promotion of "Bystander Revolution" as part of a bully cessation program, where adolescents are empowered to stop and report a bullying event.
- ii. Awareness programmes for cyberbullying should now focus on the perpetrator, as there are already approaches to manage victims of cyberbully.
- iii. In schools, life skills education and the implementation of programmes to strengthen the communication between adolescents and teachers.
- iv. The overall approach to cyber safety emphasises media watch, written policies and laws to control media contents (violence acts and pornography) and enhancing tele-health and digitalization for accessibility to get help.

### 3.12.7 References

1. WHO. (2022) Fact sheet: Adolescent and young adult health
2. Garstang, J. et al. (2020) Effect of COVID-19 lockdown on child protection medical assessments: A retrospective observational study in Birmingham, UK. *BMJ Open* 10, 1–6
3. Vaillancourt, T. et al. (2021) School bullying before and during COVID-19: Results from a population based randomized design. *Aggress. Behav.* 47, 557–569



Table 3.12.1: Prevalence of involvement in violence at least once in the past 12 months among adolescents in WP Labuan, 2022

Socio-demographic characteristics	Having been physically attacked at least once				Involvement in physical fight at least once			
	Unweighted count	Estimated population	Prevalence (%)	95 % CI Lower Upper	Unweighted count	Estimated population	Prevalence (%)	95 % CI Lower Upper
<b>WP LABUAN</b>	350	1131	17.4	15.07 20.02	409	1333	20.5	18.89 22.29
<b>Sex</b>								
Male	171	551	17.4	13.86 21.63	220	718	22.7	19.02 26.89
Female	179	580	17.4	14.42 20.86	189	615	18.5	15.49 21.88
<b>Form</b>								
Form 1	101	354	24.2	20.23 28.60	103	366	25.1	19.63 31.39
Form 2	73	252	16.9	11.49 24.16	106	374	25.1	20.44 30.44
Form 3	67	231	17.5	12.39 24.00	71	246	18.6	14.65 23.32
Form 4	57	154	13.1	7.02 23.26	76	205	17.5	8.99 31.26
Form 5	52	141	13.4	9.66 18.26	53	143	13.6	9.12 19.87
<b>Ethnicity</b>								
Malay	152	481	16.5	12.64 21.35	188	606	20.8	18.24 23.67
Chinese	32	108	13.8	8.85 21.02	38	131	16.8	12.32 22.44
Indian	1	-	-	- -	1	-	-	- -
Bumiputera Sabah	103	334	18.4	14.67 22.77	115	377	20.8	17.10 25.04
Bumiputera Sarawak	12	39	15.9	8.60 27.44	10	-	-	- -
Others	50	165	22.7	19.20 26.53	57	181	25.0	18.89 32.20

- Prevalence with high RSE, not reported

**Table 3.12.2: Prevalence of had serious injury at least once in the past 12 months among adolescents in WP Labuan, 2022**

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
<b>WP LABUAN</b>	406	1310	20.2	18.57	21.87
<b>Sex</b>					
Male	215	689	21.8	18.63	25.24
Female	191	622	18.7	16.71	20.79
<b>Form</b>					
Form 1	107	375	25.6	19.82	32.47
Form 2	93	319	21.4	16.77	26.94
Form 3	79	274	20.7	16.56	25.51
Form 4	64	177	15.1	8.40	25.64
Form 5	63	166	15.8	11.89	20.66
<b>Ethnicity</b>					
Malay	195	613	21.1	18.95	23.35
Chinese	41	138	17.7	11.39	26.45
Indian	0	-	-	-	-
Bumiputera Sabah	105	346	19.0	15.81	22.76
Bumiputera Sarawak	16	53	21.3	10.68	38.00
Others	49	161	22.1	15.35	30.85

- Prevalence with high RSE, not reported

**Table 3.12.3: Major cause of the most serious injury sustained in the past 12 months among adolescents who were injured in WP Labuan, 2022**

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
In a motor vehicle accident or hit by a motor vehicle	46	-	-	-	-
Fell	127	411	7.0	5.80	8.41
Something fell or hit him/her	20	65	1.1	0.79	1.54
Attacked of abused or fighting with someone	10	-	-	-	-
In a fire or too near a flame or something hot	4	-	-	-	-
Inhaled or swallowed something bad	6	20	0.3	0.21	0.52

- Prevalence with high RSE, not reported

Table 3.12.4: Prevalence of being abused at least once in the past 30 days among adolescents in WP Labuan, 2022

Socio-demographic characteristics	Physical abuse at home at least once				Verbal abuse at home at least once			
	Unweighted count	Estimated population	Prevalence (%)	95 % CI Lower      Upper	Unweighted count	Estimated population	Prevalence (%)	95 % CI Lower      Upper
<b>WP LABUAN</b>	194	644	9.9	7.84      12.49	954	3052	47.0	40.71      53.35
<b>Sex</b>								
Male	76	262	8.3	5.32      12.70	313	986	31.2	28.21      34.26
Female	118	382	11.5	7.66      16.86	641	2066	62.0	58.12      65.77
<b>Form</b>								
Form 1	73	255	17.4	10.65      27.25	186	651	44.5	35.71      53.63
Form 2	49	161	10.8	6.18      18.27	210	709	47.6	39.13      56.23
Form 3	37	126	9.5	5.77      15.23	185	628	47.5	34.40      60.89
Form 4	26	78	6.7	3.41      12.65	200	580	49.6	40.48      58.67
Form 5	9	25	2.4	1.32      4.27	173	484	46.1	31.40      61.47
<b>Ethnicity</b>								
Malay	88	285	9.8	7.14      13.34	411	1289	44.3	36.95      51.86
Chinese	26	92	11.9	8.15      16.96	81	273	35.0	26.82      44.18
Indian	1	-	-	-      -	2	-	-	-      -
Bumiputera Sabah	52	175	9.6	6.92      13.21	308	993	54.7	45.18      63.83
Bumiputera Sarawak	8	-	-	-      -	44	146	58.9	40.59      74.99
Others	19	62	8.5	5.14      13.85	108	345	47.5	37.68      57.55

- Prevalence with high RSE, not reported

**Table 3.12.5: Prevalence of experience in being bullied at least once in the past 30 days among adolescents in WP Labuan, 2022**

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
<b>WP LABUAN</b>	197	645	9.9	8.28	11.87
<b>Sex</b>					
Male	84	273	8.6	6.08	12.08
Female	113	372	11.2	8.48	14.60
<b>Form</b>					
Form 1	71	246	16.8	11.69	23.57
Form 2	44	148	10.0	6.57	14.82
Form 3	40	138	10.4	6.73	15.73
Form 4	27	-	-	-	-
Form 5	15	40	3.8	2.30	6.32
<b>Ethnicity</b>					
Malay	83	267	9.2	7.22	11.61
Chinese	17	-	-	-	-
Indian	3	-	-	-	-
Bumiputera Sabah	54	177	9.7	6.59	14.13
Bumiputera Sarawak	13	45	18.3	12.95	25.09
Others	27	88	12.1	8.57	16.87

- Prevalence with high RSE, not reported

**Table 3.12.6: Most common ways of being bullied at least once in the past 30 days among adolescents in WP Labuan, 2022**

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
Hit, kicked, pushed, shoved around or locked indoor	12	39	0.6	0.36	1.14
Made fun of race, nationality or color	24	77	1.3	0.87	1.79
Made fun because of religion	3	-	-	-	-
Made fun with sexual jokes, comments of gestures	27	88	1.4	0.92	2.26
Left out activities on purpose of completely ignored	13	-	-	-	-
Made fun of how body or face looks	37	121	2.0	1.56	2.51

- Prevalence with high RSE, not reported

**Table 3.12.7: Prevalence of involvement in cyberbullying activities (perpetrator) a few times within a year or more among adolescents in WP Labuan, 2022**

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
<b>WP LABUAN</b>	437	1376	21.2	17.89	24.89
<b>Sex</b>					
Male	263	818	25.9	21.40	30.88
Female	174	557	16.7	14.18	19.62
<b>Form</b>					
Form 1	82	291	19.9	12.62	29.94
Form 2	88	309	20.7	15.12	27.77
Form 3	74	263	19.8	16.28	23.94
Form 4	103	271	23.1	12.93	37.84
Form 5	90	242	23.1	15.54	32.88
<b>Ethnicity</b>					
Malay	210	646	22.2	18.26	26.73
Chinese	37	125	16.0	9.77	25.20
Indian	0	-	-	-	-
Bumiputera Sabah	118	384	21.1	16.48	26.68
Bumiputera Sarawak	12	-	-	-	-
Others	60	185	25.5	15.49	38.91

- Prevalence with high RSE, not reported

**Table 3.12.8: Most common ways of involvement in cyberbullying activities (perpetrator) a few times within a year or more among adolescents in WP Labuan, 2022**

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
Ever made rude comments to anyone online	267	840	12.9	10.06	16.47
Ever sent or posted other's embarrassing photos online	101	313	4.8	3.17	7.25
Ever spread rumours about someone online	154	480	7.4	5.79	9.41
Ever made threatening comments to hurt someone online	31	99	1.5	0.96	2.39
Ever asked someone to talk about sex online	58	186	2.9	1.86	4.39
Ever asked someone to do something sexual online	23	-	-	-	-

- Prevalence with high RSE, not reported

### 3.13 Adolescents' Perspectives on the Impact of COVID-19 on their families

Contributors: S Maria Awaluddin, Lim Kuang Kuay, Noor Syaqlah Shawaluddin, Tuan Mohd Amin Tuan Lah, Maznieda Mahjom, Noor Ani Ahmad, Saidatul Norbaya Buang, Nik Rubiah Nik Abdul Rashid.

#### 3.13.1 Introduction

The COVID-19 pandemic has impacted adolescents in many aspects, such as their developmental milestones and well-being, even though they are less affected by the COVID-19 disease<sup>1</sup>. Adolescents may have many positive and negative perspectives towards the government restrictions on gathering and outdoor activities, which will affect their mental health status<sup>2</sup>.

#### 3.13.2 Objectives

To determine the prevalence of adolescents, reported that:

- i. Parents lost their jobs due to the COVID-19 pandemic.
- ii. Their family has to cut their expenses due to the COVID-19 pandemic.
- iii. Their family needs to move to a less expensive rental house due to the COVID-19 pandemic
- iv. Their family had to sell properties due to the COVID-19 pandemic
- v. Family relationships became strained due to the COVID-19 pandemic
- vi. Family had no changes due to the COVID-19 pandemic
- vii. Their family ever been infected with COVID-19

#### 3.13.3 Variable definitions

- **Parents lost job:** the adolescent answered option "Yes" to the statement My parent/s lost his/her/their job.
- **Family has to cut their expenses:** the adolescent answered option "Yes" to the statement "Our family has to cut our expenses".
- **Family has moved to a less expensive rental house:** the adolescent answered option "Yes" to the statement My parent(s) lost his/her/their job.
- **Family had to sell properties:** the adolescent answered option "Yes" to the statement "Our family has to sell properties".
- **Family relationships became strain:** the adolescent answered option "Yes" to the statement family relationships became strained/not close.
- **Family had no changes:** adolescent who answered "No" for each subquestion of 6(a), 6(b), 6(c), 6(d), 6(e) and 6(f).
- **Family ever been infected with COVID-19:** the adolescent answered option "Yes" to the question "Has your family ever been infected with COVID-19?".

#### 3.13.4 Findings

The prevalence of adolescents in Wilayah Persekutuan Labuan reported that parents lost their job due to the COVID-19 pandemic was 9.8% (95% CI: 7.39, 12.90). Adolescents also reported that their family had to cut their expenses, needed to move to a less expensive rental house and had to sell properties, 43.2% (95% CI: 36.02, 50.75), 4.9% (95% CI: 3.42, 7.01) and 3.3% (95% CI: 2.45, 4.46), respectively. The prevalence of adolescents reported that family relationships became strained was 9.8% (95% CI: 8.78, 11.00). About 49.8% (95% CI: 43.40, 56.22) of adolescents reported that their family had no negative impact during the pandemic COVID-19. In terms of adolescents reporting family members ever been infected with COVID-19, the prevalence was 52.3% (95% CI: 47.87, 56.63). (Table 3.13.1).

#### 3.13.5 Discussion / Conclusion

About half of the adolescents reported that their family had experienced negative consequences following the pandemic COVID-19 including family members ever been infected with COVID-19. Identification of the family who experienced a negative impact due to the COVID-19 pandemic can prevent further mental health problems among the affected population.

#### 3.13.6 Recommendations

Pandemic COVID-19 caused a serious impact on the marginal of the family unit in terms of economic sustainability and parent-adolescent relationships. Therefore, the recommendations are:

- i. To work up on the emergency financial support system in reaching the affected family
- ii. To strengthen community support such as fundraising, food donation and emergency shelter
- iii. To spread awareness on Covid-19 prevention at home to prevent further disease spread
- iv. To encourage adolescents to reach out for help in any difficult situation.

#### 3.13.7 References

1. Rao N, Fisher PA. The impact of the COVID-19 pandemic on child and adolescent development around the world. *Child Dev.* 2021;92(5): e738–48
2. Panchal U, Salazar de Pablo G, Franco M, Moreno C, Parellada M, Arango C, et al. The impact of COVID-19 lockdown on child and adolescent mental health: systematic review. *Eur Child Adolesc Psychiatry* [Internet]. 2021;(0123456789). Available from: <https://doi.org/10.1007/s00787-021-01856-w>

**Table 3.13.1: Adolescents’ perspectives on the impact of COVID-19 pandemic to their family: Self-reported findings from the adolescents in WP Labuan, 2022**

Categories of COVID-19 impact	Unweighted count	Estimated population	Prevalence (%)	95 % CI	
				Lower	Upper
Parents lost job	200	637	9.8	7.39	12.90
Family had to cut their expenses	891	2805	43.2	36.02	50.75
Family needed to move to less expensive rental house	99	319	4.9	3.42	7.01
Family had to sell properties	66	215	3.3	2.45	4.46
Family relationships became strain	198	638	9.8	8.78	11.00
Family had no changes	1004	3233	49.8	43.40	56.22

## APPENDICIES

### Appendix 1: Members of Steering Committee NHMS 2019-2022

1. Director General of Health
2. Deputy Director General of Health (Research & Technical Support)
3. Deputy Director General of Health (Public Health)
4. Deputy Director General of Health (Medical)
5. Principal Director, Oral Health Programme
6. Principal Director, Pharmaceutical Services
7. Principal Director, Food Safety and Quality Division
8. Director, Medical Development Division
9. Director, Planning Division
10. Director, Health Education Division
11. Director, Disease Control Division
12. Director, Family Health Development Division
13. Director, Nutrition Division
14. Representative of State Directors
15. Director, Institute for Public Health
16. Dean Faculty of Medicine, University of Malaya
17. Dean Faculty of Medicine, National University of Malaysia
18. Principal Investigator, NHMS

### Appendix 2: Terms of reference for NHMS 2022 Steering Committee

1. To approve the objectives and scopes of NHMS 2019 - 2022
2. To facilitate inter and intra sectorial collaboration
3. To monitor the implementation of NHMS 2019 - 2022
4. To facilitate the utilization of the NHMS 2019 - 2022 findings

### Appendix 3: List of members of Central Coordinating Committee, NHMS 2022

1. Dr. Noor Ani Ahmad, Director of Institute for Public Health
2. Mr. Lim Kuang Kuay, Principal Investigator of Adolescent Health Survey
3. Dr. Muhammad Fadhli Mohd Yusoff, Method And Statistic
4. Ms. Hamizatul Akmal Abd Hamid, Data Manager
5. Dr. Shubash Shander Ganapathy, Central Field Supervisor of Negeri Sembilan, Melaka & Johor
6. Dr. Ahmad Ali Hj Zainuddin, Central Field Supervisor of Kedah, Perlis, Pulau Pinang & Perak
7. Dr. S Maria Awaluddin, Central Field Supervisor of Kelantan, Terengganu & Pahang
8. Dr. Maznieda Mahjom, Central Field Supervisor of Selangor, Kuala Lumpur & Wilayah Putrajaya
9. Dr. Mohd Shaiful Azlan Kassim, Central Field Supervisor of Sabah, Labuan & Sarawak
10. Dr. Tan Lee Ann, Data Processing & Quality
11. Ms Noor Syaqilah Shawaluddin, Logistic Support
12. Ms. Nashrah Adilah Ismail, Project Manager



**Appendix 4: Terms of Reference for NHMS 2022 Central Coordinating Team (CCT)**

No	Team	Duties	Officers
1	Project Management and Finance	<ul style="list-style-type: none"> <li>• Work closely with recruitment group for employment of temporary Research Assistant</li> <li>• Prepare Questionnaires manual, Data collection manual</li> <li>• Meeting with research team members, and stakeholders</li> <li>• Planning for data collection training</li> <li>• Prepare security cards/name tags for research team Arrangement for advanced payment for survey research teams</li> <li>• Process claims of Field Supervisors</li> <li>• Monitor the expenditure/budget</li> </ul>	<p>Mr. Lim Kuang Kuay                      Dr. S. Maria Awaluddin                      Ms. Nashrah Adilah Ismail Ms.                      Nurul Amalina Yusof                      Mr. Muhammad Safuan Suhaimi                      Mr. Mohamad Shafiq Abd Basid</p>
2	Method and Data Analysis	<p><b>Before Data Collection</b></p> <ul style="list-style-type: none"> <li>• Calculate the sample size</li> <li>• Determine the sample distribution by state</li> <li>• Selection of schools samples for each state</li> <li>• Selection of class samples from the selected school</li> <li>• Prepare unique ID for the selected schools and classes</li> <li>• Check module cover and dummy table prepared by key module</li> </ul> <p><b>During Data Collection</b></p> <ul style="list-style-type: none"> <li>• Monitoring the quality of data received from data processing team</li> <li>• Conducting daily data cleaning</li> <li>• Merge the clean dataset</li> <li>• Updating the monitoring board for data processing and data quality during CCT meeting</li> <li>• Analyst the estimate prevalence for each module during data collection</li> </ul> <p><b>After Data Collection</b></p> <ul style="list-style-type: none"> <li>• Check syntax analysis to ensure the analysis meet the module objectives</li> <li>• Prepare sampling weight for complex sample analysis</li> <li>• Check the table analysis for technical report</li> <li>• Prepare final database</li> <li>• Prepare data dictionary for reference</li> </ul>	<p>Ms. Hamizatul Akmal Abd Hamid                      Dr. Muhammad Fadhli Mohd Yusoff                      Ms. Nur Syahirah Ibrahim</p>
3	Data Processing and Quality	<ul style="list-style-type: none"> <li>• Setting up data processing facility Development of directory of variables database</li> <li>• Development of quality control (QC) manual for data processing Specify data structure for data processing and data output requirement</li> <li>• Responsible for data entry and data cleaning</li> <li>• Monitoring and evaluation of QC performance for data processing Maintenance of the scanning machine</li> <li>• Daily back up for databases</li> </ul>	<p>Dr. Tan Lee Ann                      Ms. Nurul Haniyah Roslan Ms.                      Nur Faraeein Zainal Abidin                      Ms. Azlin Awatif Mohd Amir Hamzah</p>

No	Team	Duties	Officers
4	Central Field Supervisors	<p><b>Before Data Collection</b></p> <ul style="list-style-type: none"> <li>Central Field Supervisors are expected to prepare for the initiation of data collection. The preparation tasks include: <ul style="list-style-type: none"> <li>Conduct meeting with State Education Office, School Principals, Teacher in-charged for the selected schools.</li> <li>To ensure adequate logistic support for the data collection and liaise with the District Education Office, District Health Office and other relevant departments to ensure that: <ul style="list-style-type: none"> <li>Human resources are available: Field Supervisors, Team leaders, Research Assistants and drivers.</li> <li>Manage transport: Vehicles</li> <li>Manage survey instruments and relevant form</li> <li>Manage lodging for data collectors</li> </ul> </li> </ul> </li> </ul> <p><b>During Data Collection</b></p> <ul style="list-style-type: none"> <li>Gather feedback from the field on the data collection status and problems related to logistics.</li> <li>Visit the field to help data collectors solve the problem if necessary.</li> <li>To ensure all data collection monitoring forms have been received on time.</li> <li>To ensure bundle from field received by the Operation Centre by hand and by post.</li> <li>Updating the monitoring board for state achievement and attending CCT meeting.</li> </ul>	<p>Dr. Ahmad Ali Zainuddin  Dr. Maznieda Mahjom  Dr. Mohd Shaiful Azlan Kassim  Dr. S Maria Awaluddin  Dr. Shubash Shander Ganapathy</p>
5	Operation Centre	<ul style="list-style-type: none"> <li>Arrange date and place of meeting</li> <li>Prepare and circulate briefing materials</li> <li>Prepare and circulate minutes of CCT meeting</li> <li>Prepare letters of appointment for Central Field Supervisors, Field Supervisors and data collectors</li> <li>Prepare advertisement material for recruitment of data collectors</li> <li>Prepare letters of notifications for data collections</li> <li>Prepare manuals for field Supervisors and data collectors</li> <li>Develop a system/format and monitor the distribution of materials/ equipment for field work</li> </ul>	<p>Mr. Lim Kuang Kuay  Ms. Nashrah Adilah Ismail</p>

## Appendix 5: List of Research Team Members, NHMS 2022

### Alcohol Use

1. Dr. Rusdi Abd Rashid
2. Dr. Norli Abdul Jabbar
3. Mr. Faizul Akmal Abdul Rahim
4. Ms. Hamizatul Akmal Abd Hamid
5. Ms. Halizah Mat Rifin
6. Ms. Hasimah Ismail
7. Mr. Mohd Hatta Abdul Mutalip
8. Dr. Muhammad Fadhli Mohd Yusoff
9. Dr. Thamil Arasu Saminathan
10. Dr. Tania Gayle Robert
11. Dr. Chong Zhuo Lin

### Dietary Behaviours

1. Dr. Ahmad Ali Zainuddin
2. Ms. Ainan Nasrina Ismail
3. Ms. Teh Wai Siew
4. Dr. Lai Wai Kent
5. Dr. Suhaila Abdul Ghaffar
6. Mr. Azli Bin Baharudin@Shaharudin
7. Mr. Chong Chean Tat
8. Ms. Lalitha Palaniveloo
9. Mr. Muhammad Faiz Mohd Hisham
10. Ms. Munawara Pardi
11. Dr. Norsyamliana Che Abdul Rahim
12. Ms. Nurul Huda Ibrahim
13. Ms. Siti Adibah Ab. Halim
14. Ms. Syafinaz Mohd Sallehuddin

### Nutritional Status

1. Ms. Ainan Nasrina Ismail
2. Dr. Ahmad Ali Zainuddin
3. Ms. Lalitha Palaniveloo
4. Mr. Khairul Hasnan Amali
5. Ms. Siti Adibah Ab. Halim

### Drug Use

1. Dr. Mohamad Salleh Abdul Ghani
2. Dr. Norli Abdul Jabbar
3. Dr. Rushidi Abd Rashid
4. Dr. Thamil Arasu Saminathan
5. Dr. Maznieda Mahjom
6. Ms. Hasimah Ismail
7. Ms. Hamizatul Akmal Abd Hamid
8. Dr. Muhammad Fadhli Mohd Yusoff
9. Mr. Mohd Haniff Bistari
10. Dr. Halizah Mat Rifin
11. Dr. Tania Gayle Rober

### Hygiene (Including Oral Health)

1. Dr. Fazila Haryati
2. Ms. Rafidah Ali
3. Dr. Chan Yee Mang
4. Mr. Mohd Hatta Abdul Mutalip
5. Dr. Nik Adilah Shahein
6. Ms. Norzawati Yoep
7. Dr. Annapurny Venkiteswaran
8. Dr. Nurulasmak Mohamed
9. Dr. Nik Daliana Nik Farid
10. Dr. Saidatul Norbaya Buang

### Mental Health Problems

1. Dr. Nurashikin Ibrahim
2. Dr. Nor Rahidah Abdul Rahim
3. Dr. Noor Raihan Khamal
4. Dr. Noor Ani Ahmad,
5. Dr. Sherina Mohd Sidek
6. Ms. Norhafizah Sahril
7. Dr. Chan Yee Mang
8. Dr. Kishwen Kanna Yoga Ratnam
9. Mr. Mohd Ruhaizie Riyadzi
10. Mr. Mohd Haniff Bistari
11. Dr. Muhammad Azri Adam Adnan
12. Dr. Muhamad Khairul Nazrin Khalil
13. Dr. Mohd Shaiful Azlan Kassim
14. Mr. Mohamad Aznuddin Abd Razak
15. Ms. Nur Hidayatun Fadhilah Mohd Nor
16. Mr. Sheikh Shafizal Sheikh Iلمان

### Physical Activity

1. Dr. Hazizi Abu Saad
2. Dr. Mohd Azahadi Omar
3. Ms. Nur Hidayatun Fadhilah Mohd Nor
4. Dr. Muhammad Solihin Rezali
5. Dr. Affendi Isa
6. Ms. Siti Balkhis Shafie
7. Mr. Lim Kuang Kuay
8. Mr. Mohamad Aznuddin Abd Razak
9. Dr. Mohd Shaiful Azlan Kassim
10. Mr. Azli Baharudin@Shaharudin
11. Mr. Mohd Hairmansah Mohd Shah
12. Ms. Nor'Ain Ab Wahab
13. Ms. Norliza Shamsuddin
14. Ms. Nazirah Alias
15. Ms. Nurul Haniyah Rosslan

### **Protective Factors**

1. Dr. Nik Rubiah Nik Abdul Rashid
2. Dr. Nik Daliana Nik Farid
3. Dr. Zamzaireen Zainal Abidin
4. Ms. Nazirah Alias
5. Ms. Eida Nurhadzira Muhammad
6. Ms. Filza Noor Asari
7. Mr. Faizul Akmal Abdul Rahim
8. Dr. Tan Lee Ann
9. Dr. S Maria Awaluddin
10. Dr. Khaw Wan-Fei
11. Mr. Mohd Amierul Fikri Mahmud
12. Mr. Mohd Farihan Md Yatim
13. Dr. Nur Hamizah Nasaruddin

### **Sexual Behaviour that contribute to HIV infection, other STI and unintended pregnancy**

1. Dr. Anita Suleiman
2. Dr. Nik Rubiah Nik Abdul Rashid
3. Dr. Chong Zhuo Lin
4. Dr. Fatin Athira Tahir
5. Dr. Mazliza Ramly
6. Dr. Maznieda Mahjom
7. Dr. Nik Adilah Shahein
8. Dr. S Maria Awaluddin
9. Dr. Noor Aliza Lodz
10. Dr. Amal Shamsudin

### **Tobacco Use**

1. Dr. Noraryana Hassan
2. Dr. Norliana Ismail
3. Dr. Muhammad Hairul Nizam Abd Hamid
4. Ms. Ummi Nadiah Yusoff
5. Dr. Nizam Baharom
6. Mr. Lim Kuang Hock
7. Mr. Mohd Ruhaizie Riyadzi
8. Dr. Muhammad Fadhli Mohd Yusoff
9. Dr. Thamil Arasu Saminathan
10. Dr. Tania Galye Robert Lourdes
11. Dr. Halizah Mat Rifin
12. Ms. Hamizatul Akmal Abd Hamid
13. Ms. Hasimah Ismail
14. Dr. Wan Kim Sui
15. Dr. Kishwen Kanna Yoga Ratnam

### **Violence and Unintentional Injury**

1. Ms. Hamizatul Akmal Abd Hamid
2. Dr. Tan Lee Ann
3. Dr. Nor Rahidah Abd Rahim
4. Dr. Noor Raihan Khamal
5. Mr. Mohd Hazrin Hasim@Hashim
6. Ms. Nur Faraeein Zainal Abidin
7. Dr. Noor Suraya Muhamad
8. Dr. Shubash Shander Ganapathy
9. Mr. Muhammad Hanafi Bakri

### **Adolescents' Perspective on the Impact of COVID-19 to their family**

1. Dr. S Maria Awaluddin
2. Mr. Lim Kuang Kuay
3. Ms. Noor Syaqlah Shawaluddin
4. Mr. Tuan Mohd Amin Tuan Lah
5. Dr. Maznieda Mahjom
6. Dr. Noor Ani Ahmad
7. Dr. Saidatul Norbaya Buang
8. Dr. Nik Rubiah Nik Abdul Rashid

**Appendix 6: List of Data Collection Teams, NHMS 2022**

**JOHOR**

**Field Supervisor**

Dr. Lai Wai Kent

**Drivers**

1. Mr. Muhammad Azraei Alias
2. Mr. Mohammad Nazrin Nazmuding

**Research Assistants**

1. Ms. Salsabeela Mohd Ariff
2. Ms. Nurfatin Syazwana Ayob
3. Ms. Raja Nur Fatin Ainsyah Raja Omar
4. Ms. Nor Diana Zulkefli
5. Mr. Mohammad Luqman Abdul Aziz
6. Ms. Siti Noorul Nadhirah Zamrus

**KEDAH**

**Field Supervisor**

Mrs. Lalitha Palaniveloo

**Drivers**

1. Mr. Muhammad Shahrul Arieff Shahrudin
2. Mr. Mohamad Najmi Shahrin

**Research Assistants**

1. Ms. Nur Liyana Rosle
2. Mr. Muhammad Iqbal Mat Rosdi
3. Ms. Siti Nur Adibah Zainudin
4. Ms. Nur Hawanis Hashim
5. Mr. Muhammad Zaquan Mohamad Zamri
6. Ms. Noor Fazira Mhd Sofbri

**KELANTAN**

**Field Supervisor**

Dr. Norsyamalina Che Abdul Rahim

**Drivers**

1. Mr. Muhamad Sahasrizan Samat
2. Mr. Muhamad Izzat Amir Mohd Nasir

**Research Assistants**

1. Mr. Muhammad 'Izzuddin Che Ismail
2. Mr. Mohamad Azli Che Daud
3. Ms. Wan Anisa Rodzlan Hasani
4. Mr. Muhammad 'Izzuddin Che Ismail
5. Ms. Nurul Farhani Faizol
6. Ms. Siti Hajar Ishak

**MELAKA**

**Field Supervisor**

Ms. Eida Nurhadzira Muhammad

**Drivers**

1. Ms. Siti Zulaikha Yahya
2. Ms. Puteri Nurdhiyana Othman

**Research Assistants**

1. Ms. Erma Safwan Erison
2. Ms. Nur Aishah Solihin Mohmad Nezan
3. Ms. Siti Normah Abdul Manan
4. Ms. Najihah Md Din
5. Ms. Nur Anis Syafiqqa Zulkefli
6. Ms. Fairuz Mohd Hashim

**NEGERI SEMBILAN**

**Field Supervisor**

Mr. Jayvikramjit Singh Manjit Singh

**Drivers**

1. Mr. Zakaria Mohammad
2. Mr. Gabriel Jatum

**Research Assistants**

1. Ms. Norsahira Kamarudin
2. Mr. Mohamad Pauzan Razali
3. Ms. Norhayati Kamarudin
4. Ms. Nurul Syuhada Samsuddin
5. Ms. Siti Aisyah Ibrahim
6. Ms. Izzati Wan Azelee

**PAHANG**

**Field Supervisor**

Mr. Sheikh Shafizal Sheikh Ilman

**Drivers**

1. Mr. Muhammad Ruzaini Ahmad Amri
2. Mr. Ihsan Hashim

**Research Assistants**

1. Ms. Norhakimah Md Din
2. Mr. Harizamharizal Syafrizal
3. Ms. Norhidayah Abdul Majid
4. Ms. Nur Aina Amira Zailani
5. Ms. Geerthana A/P R. Ravichandiran
6. Mr. Muhamad Firdaus Paizol

**PULAU PINANG****Field Supervisor**

Ms. Rafidah Ali

**Drivers**

1. Mr. Muhammad Arif Misra
2. Mr. Muhammad Syauqi Adrus

**Research Assistants**

1. Mrs. Eng Gaik Sim
2. Mr. Neoh Choo Loa
3. Mr. Mohammad Hasrizal Hassan
4. Mr. Tan Jun Xian
5. Mr. Muhammad Amin Sabri
6. Ms. Nurnabilah Afrina Azami

**PERAK****Field Supervisor**

Dr. Halizah Mat Riffin

**Drivers**

1. Mr. Muhammad Raidillah Che Ab. Rahim
2. Mr. Muhamad Syawal Azim Mohd Hisham

**Research Assistants**

1. Ms. Azieda Abu Bakar
2. Ms. Zawahir Ngah Said
3. Ms. Erma Natasa Norhan
4. Ms. Amni Zulaika Ahmad Azmi
5. Ms. Hasziefatul Affidah Hasnan
6. Mr. Amirul Amin Mohamed Tarmizi

**PERLIS****Field Supervisor**

Dr. Suhaila Abdul Ghaffar

**Drivers**

1. Mr. Mohammad Amiruddin Kamarunzaman
2. Mr. Mohd Aizam Zahid

**Research Assistants**

1. Ms. Ainul Mardhiah Pakhurrrazi
2. Ms. Nur Syuhada Zahid
3. Ms. Fairuz Tasnim Shaffie
4. Ms. Nor Najihah Muslim
5. Ms. Jaizah Jamil
6. Ms. Noor Faralina Izzati Kamarunzaman

**SELANGOR****Field Supervisor**

Ms. Nazirah Alias

**Drivers**

1. Mr. Hezri Izuan Ahmad Termizi
2. Mr. Muhammad Izzat Mat Yusoff

**Research Assistants**

1. Ms. Nurul Atiqah Mat Yusoff
2. Ms. Rabi'ahatul Assuhadah Mohd Rafa'ai
3. Ms. Fatini Abd Rahman
4. Mr. Muhammad Azrol Mohd Rozi
5. Ms. Noor Aiman Afaf Afiffudden
6. Ms. Nurul Ashikin Nosarodin

**TERENGGANU****Field Supervisor**Dr. Fazila Haryati Ahmad  
Mr. Mohd Ruhaizie Riyadzi**Drivers**

1. Mr. Muhammad Afif Bani Yami
2. Mr. Muhammad Najmi Alif Muda

**Research Assistants**

1. Mr. Alif Amirul Ikhwan Hussin
2. Ms. Nur Alis Nadia Azman
3. Ms. Nurul Shafiqah Kusno
4. Ms. Siti Nur Sharmiela Ayob
5. Ms. Madhiah Che Man
6. Ms. Nur Atiqah Hazwani Mohammed

**SABAH****Field Supervisor**

Ms. Nur Faraeein Zainal Abidin

**Drivers**

1. Mr. Steve Glantdenventur E Benjamin
2. Mr. Javixsen James
3. Mr. Mohd Jazlan Harith Abdul Razak

**Research Assistants**

1. Mr. Joel Sonny Saimin
2. Ms. Haslinda Hasan
3. Mr. Wan Misly Kindon
4. Ms. Nur Maisarah Maksud
5. Mr. Mohd Hafizan Sani
6. Mr. Mohd Aldy Abdul Razak
7. Ms. Marini Juanah Mantigang
8. Ms. Fyrah James
9. Ms. Lovera Karera Kalaka

**SARAWAK****Field Supervisor**Dr. Khaw Wan Hei  
Mr. Mohd Hairmanshah Mohd Shah**Drivers**

1. Mr. Mohammed Hefalani Mohd Azman
2. Mr. Wilkinson Anak Welling
3. Mr. Afiq Fakrul Ismail

**Research Assistants**

1. Mr. Daniel Sia Pong Chai
2. Ms. Nurul Afifah Nasir
3. Mr. Mugang Anak Japar
4. Mr. Fabian Anak Mathew
5. Ms. Aelsa Anak Anthony
6. Ms. Christina Sie Fang Yun
7. Ms. Happilyn Anak Li
8. Mr. Mohammad Hasnol Abd Halim
9. Mr. Nazran Bazlan Nawi

## KUALA LUMPUR

### Field Supervisor

Dr. Nur Hamizah Nasaruddin

### Drivers

1. Mr. Muhammad Muazzam Abdul Rahman
2. Mr. Adib Iman Osman

### Research Assistants

1. Mr. Amirah Ali
2. Mr. Muhammad Yusri Abdullah
3. Ms. Nur Amirah Alias
4. Mr. Nurulnatasha Jumali
5. Ms. Noor Hasnieza Ahmad
6. Mr. Muhammad Naim Ismail

## WP LABUAN

### Field Supervisor

Dr. Muhammad Azri Adam Adnan

### Driver

1. Mr. Niveno Eldo Sonny Mat

### Research Assistants

1. Ms. Mawarsari Said
2. Mr. Jeldy Galoh
3. Ms. Nor Syafina Gorganius
4. Mr. Ignasius Cartilo Taimin
5. Ms. Norfazirah Amlan
6. Ms. Noor Azni Adzmain

## WP PUTRAJAYA

### Field Supervisor

Ms. Syafinaz Mohd Sallehuddin

### Drivers

1. Mr. Mohd Sanusi Aziz
2. Mr. Muhammad Asyraf Jasri

### Research Assistants

1. Ms. Aini Farina Mohd Zamri
2. Ms. Nurul Atikah Mohd Rozi
3. Mr. Nurlis Yunarlis
4. Ms. Azizah Nurfauziah Jafri
5. Ms. Rohana Saharudin









KEMENTERIAN KESIHATAN MALAYSIA



KEMENTERIAN PENDIDIKAN MALAYSIA

ISBN 978-967-5340-75-8



9 789675 340758